

Comentarios sobre la TCC en el espacio de coordenadas

Remarks on QFT in the Coordinate Space

A. MUCH¹

Recibido: 7 de enero de 2018/ Aceptado: 17 de abril de 2018

¹Centro de Ciencias Matemáticas, Universidad Nacional Autónoma de México, C.P. 58190, Morelia, Michoacán México, email: amuch@matmor.unam.mx

Agradecimientos

El autor quisiera agradecer al Prof. K. Sibold por dar comienzo a las preguntas conceptuales profundas de este trabajo. Además, quisieramos agradecer al Prof. K. Sibold, Prof. Yuri Bonder y S. Pottel por muchas discusiones fructíferas durante las diferentes etapas de este artículo. Las correcciones lingüísticas hechas por el Dr. Z. Much son reconocidas agradecidamente.

Acknowledgment

The author would like to thank Prof. K. Sibold for initiating deep conceptual questions of this work. Moreover, we would like to thank Prof. K. Sibold, Prof. Yuri Bonder and S. Pottel for many fruitful discussions during different stages of this paper. The linguistic corrections of Dr. Z. Much are thankfully acknowledged.

Generators of the Poincaré group, for a free massive scalar field, are usually expressed in the momentum space. In this work we perform a transformation of these generators into the coordinate space. This (spatial)-position space is spanned by eigenvectors of the Newton-Wigner-Pryce operator. The motivation is twofold. First, we want to investigate the localization of a relativistic particle. Furthermore, we need a deeper understanding of the commutative spatial coordinate space in QFT, in order to investigate the non-commutative version thereof.

Los generadores del grupo de Poincaré, para un campo escalar masivo libre, se suelen expresar en el espacio de momentos. En este trabajo realizamos una transformación de estos generadores al espacio de coordenadas. Este espacio de posiciones (espaciales) es generado por eigenvectores del operador Newton-Wigner-Pryce. La motivación es doble. Primero, queremos investigar la localización de una partícula relativista. Además necesitamos un entendimiento más profundo del espacio de coordenadas espaciales conmutativo de la TCC para poder investigar su versión no conmutativa.

PALABRAS CLAVES

Fundamentos de la Teoría Cuántica de Campos, Segunda Cuantización, Operador de Coordenadas, Operador Newton-Wigner, TCC en el espacio de coordenadas, Transformación de Fourier.

KEYWORDS

Foundations of Quantum Field Theory, Second Quantization, Coordinate Operator, Newton-Wigner Operator, QFT in coordinate space, Fourier-transformation

PACS

11.10.Cd, 11.10.Nx, 11.30.Cp, 11.90.+t, 03.65.-w, 03.65.Fd, 03.70.+k, 03.30.+p

I | INTRODUCTION

Form an algebraic point of view, standard problems in quantum mechanics can be described in two spaces, the momentum and the coordinate space. These spaces are generated by two complimentary observables, which are each represented by self-adjoint operators. These operators are the momentum and

* Esta obra está bajo una licencia Creative Commons Reconocimiento - NoComercial 4.0 Internacional © ⓘ

* This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. © ⓘ

the coordinate operator. In particular, the spaces they generate consist of their respective eigenvectors. Both spaces are equally relevant for investigation and the choice usually depends on the nature of the problem.

Nevertheless, due to the complementarity, it is difficult, if not impossible, to understand quantum mechanics from both perspectives at the **same time**. In a letter from Pauli to Heisenberg (19. October 1926), there is the famous quote describing this particular problem: *„one can see the world with the p -eye or one can see the world with the x -eye, if one opens both eyes at once, one becomes crazy.“*

However, in quantum field theory (QFT) on Minkowski space the world is mostly seen through the eye of momentum space. This is partly due to the Wigner classification in terms of unitary, irreducible representations of the Poincaré group. Hence, most work is done in the momentum basis and the conjugate basis of the momentum, i.e. the spatial coordinate space is, for the most part, overlooked.

It should be noted that there exists an appropriate operator, whose eigenstates span the spatial position space. It is in literature mostly referred to as the Newton-Wigner-Pryce (NW)-operator, see (Pryce, 1948), (Newton and Wigner, 1949), (Schweber, 1961), (Jordan, 1980), (Sibold and Solard, 2009) and (Much, 2015). The importance of these states relies on the fact that they can be used to calculate the probability amplitude of finding a particle at a certain spatial-region at time t . Moreover, the second quantization of this operator allows us to calculate the probability amplitude of finding k -particles at certain spatial-positions at time t .

The motivation to perform investigations in the coordinate space, comes from the desire to understand and investigate the non-commutative version thereof. Hence, concerning investigations in QFT that assume a constant non-commutative space (without the involvement of time) the coordinate space is of great importance. In particular, it is the space where non-commutativity is introduced. Therefore, in order to understand how the Poincaré group acts or which form the generators take on non-commutative spaces, in a quantum field theoretical context, we first need to investigate the commutative framework.

Therefore, in this work we open the other eye, i.e. take a first step in this direction by expressing second-quantized quantities of the Poincaré algebra in the coordinate basis. In particular, we express generators of the respective algebras in terms of Fock space operators, which are usually written in the momentum space, and change the basis to the coordinate space. This is done by using the eigenvectors and corresponding ladder operators of the spatial coordinate operator. Moreover, we investigate how the coordinate space behaves under transformations of the Poincaré group.

It is important to point out that there is a clear distinction between the so-called configuration space (see (Schweber, 1961, Chapter 7)) and the complimentary space of the momentum, i.e. the coordinate space. In QM these two spaces are equal, however in the QFT-case these two cases are different, but related by an integral transformation.

The paper is organized as follows; Section two comprises the preliminaries, where we define the Fock space of the free scalar field and the algebra of the Fourier transformed ladder operators. The third section gives a treatment of the Newton-Wigner-Pryce operator, i.e. the respective spatial-position coordinate operator. In section four we transform the Poincaré algebra of a massive scalar field into the coordinate space. Section five investigates the transformation behavior of the coordinate space under the Poincaré group.

Conventions I.1. We use $d = n + 1$, for $n \in \mathbb{N}$ and the Greek letters are split into $\mu, \nu = 0, \dots, n$. Moreover, we use Latin letters for the spatial components which run from $1, \dots, n$ and we choose the following convention for the Minkowski scalar product of d -dimensional vectors, $a \cdot b = a_0 b^0 + a_k b^k = a_0 b^0 - \vec{a} \cdot \vec{b}$, i.e. $\eta_{jk} = \text{diag}(-1, -1, -1)$.

II | PRELIMINARIES

1 | Bosonic Fock Space and Fourier-Transformation

The $(n+1)$ -dimensional ($n \in \mathbb{N}$) relativistic Bosonic Fock space is defined in the following. Let a particle have momentum $\mathbf{p} \in \mathbb{R}^n$. Then the energy of a massive particle is $\omega_{\mathbf{p}} = +\sqrt{\mathbf{p}^2 + m^2}$. In addition, the Lorentz-invariant measure is given by $d^n \mu(\mathbf{p}) = d^n \mathbf{p} / (2\omega_{\mathbf{p}})$.

Definition The **Bosonic Fock space** $\mathcal{F}^+(\mathcal{H})$ is defined as in (Bratteli and Robinson, 1996):

$$\mathcal{F}^+(\mathcal{H}) = \bigoplus_{k=0}^{\infty} \mathcal{H}_k^+,$$

where $\mathcal{H}_0 = \mathbb{C}$ and the symmetric k -particle subspaces are given as

$$\begin{aligned} \mathcal{H}_k^+ &= \{\Psi_k : \underbrace{\mathcal{H}_1 \times \dots \times \mathcal{H}_1}_{k\text{-times}} \rightarrow \mathbb{C} \text{ symmetric} \mid \\ &\|\Psi_k\|^2 = \int d^n \mu(\mathbf{p}_1) \dots \int d^n \mu(\mathbf{p}_k) |\Psi_k(\mathbf{p}_1, \dots, \mathbf{p}_k)|^2 < \infty\}, \end{aligned}$$

with \mathcal{H}_1 being defined by

$$\mathcal{H}_1 = \{\Psi : H_m^+ \rightarrow \mathbb{C} \mid \|\Psi\|^2 = \int d^n \mu(\mathbf{p}) |\Psi(\mathbf{p})|^2 < \infty\},$$

where H_m^+ is the orbit

$$H_m^+ := \{p \in \mathbb{R}^{n+1} \mid p^2 = m^2, p_0 > 0\}.$$

The ladder operators a, a^* for this particular space are defined as follows.

Definition The covariant **particle annihilation** and **creation operators** are defined by their action on k -particle wave functions,

$$\begin{aligned} (a_c(f)\Psi)_k(\mathbf{p}_1, \dots, \mathbf{p}_k) &= \sqrt{k+1} \int d^n \mu(\mathbf{p}) \overline{f(\mathbf{p})} \Psi_{k+1}(\mathbf{p}, \mathbf{p}_1, \dots, \mathbf{p}_k) \\ (a_c(f)^*\Psi)_k(\mathbf{p}, \mathbf{p}_1, \dots, \mathbf{p}_k) &= \begin{cases} 0, & k = 0 \\ \frac{1}{\sqrt{k}} \sum_{i=1}^k f(\mathbf{p}_i) \Psi_{k-1}(\mathbf{p}_1, \dots, \mathbf{p}_{i-1}, \mathbf{p}_{i+1}, \dots, \mathbf{p}_k), & k > 0 \end{cases} \end{aligned}$$

with $f \in \mathcal{H}_1$ and $\Psi_k \in \mathcal{H}_k^+$. The commutator relations of $a_c(f), a_c(f)^*$ follow immediately and are given as

$$[a_c(f), a_c(g)^*] = (f, g) = \int d^n \mu(\mathbf{p}) \overline{f(\mathbf{p})} g(\mathbf{p}), \quad [a_c(f), a_c(g)] = 0 = [a_c(f)^*, a_c(g)^*]$$

Particle annihilation and creation operators with sharp momentum are introduced as operator valued distributions and are given by

$$a_c(f) = \int d^n \mu(\mathbf{p}) \overline{f(\mathbf{p})} a_c(\mathbf{p}), \quad a_c(f)^* = \int d^n \mu(\mathbf{p}) f(\mathbf{p}) a_c^*(\mathbf{p}),$$

where the particle annihilation and creation operators with sharp momentum satisfy the following commutator relations

$$[a_c(\mathbf{p}), a_c^*(\mathbf{q})] = 2\omega_{\mathbf{p}} \delta^n(\mathbf{p} - \mathbf{q}), \quad [a_c(\mathbf{p}), a_c(\mathbf{q})] = 0 = [a_c^*(\mathbf{p}), a_c^*(\mathbf{q})] \quad (1)$$

From a book keeping point of view, the noncovariant representation of the annihilation and creation operators given as,

$$a(\mathbf{p}) := \frac{a_c(\mathbf{p})}{\sqrt{2\omega_{\mathbf{p}}}}, \quad a^*(\mathbf{p}) := \frac{a_c^*(\mathbf{p})}{\sqrt{2\omega_{\mathbf{p}}}},$$

is easier to handle. Hence, in the following sections, when explicit calculations are performed, we use the non-covariant representation. In order to give the base change of the infinitesimal generators of the Poincaré group, in terms of the Fourier-transformed creation and annihilation operators, we first explicitly define the base change.

Definition Fourier-transformation

The base change to the Fourier-transformed creation and annihilation operators is given by,

$$a(\mathbf{p}) = (2\pi)^{-n/2} \int d^n \mathbf{x} e^{ip_k x^k} \tilde{a}(\mathbf{x}), \quad a^*(\mathbf{p}) = (2\pi)^{-n/2} \int d^n \mathbf{x} e^{-ip_k x^k} \tilde{a}^*(\mathbf{x}).$$

From the commutation relation between the two operators a and a^* we can deduce the relations for the Fourier-transformed operators,

$$\delta^n(\mathbf{p} - \mathbf{q}) = [a(\mathbf{p}), a^*(\mathbf{q})] = (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} e^{ip_k x^k} e^{-iq_k y^k} [\tilde{a}(\mathbf{x}), \tilde{a}^*(\mathbf{y})].$$

Hence the solution of the commutation relations of the coordinate space creation and annihilation operators is given, as follows

$$[\tilde{a}(\mathbf{x}), \tilde{a}^*(\mathbf{y})] = \delta^n(\mathbf{x} - \mathbf{y}).$$

The inverse transformations are given by,

$$\tilde{a}(\mathbf{x}) = (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-ip_k x^k} a(\mathbf{p}), \quad \tilde{a}^*(\mathbf{x}) = (2\pi)^{-n/2} \int d^n \mathbf{p} e^{ip_k x^k} a^*(\mathbf{p}). \quad (2)$$

This commutator could also be obtained by taking the CR's (commutator relations) of the smeared operators into account. Moreover, on eigenvectors of the coordinate operator, the Fourier-transformed creation and annihilation operators act \tilde{a} , \tilde{a}^* as ladder operators. Note that the definition of the Fourier-transformed operators does not explicitly depend on the mass.

2 | Constantly Used Integrals

In order to make this work self-contained we give the general formula for certain Fourier-transformed functions, (Gel'fand and Shilov, 1964, Chapter III, Section 2.8)

$$\begin{aligned}\tilde{p}^\lambda &= \int d^n \mathbf{p} (|\vec{p}|^2 + m^2)^\lambda \exp(-i p_k z^k) \\ &= \frac{2^{\lambda+1} (2\pi)^{\frac{1}{2}n}}{\Gamma(-\lambda)} \left(\frac{m}{|\vec{z}|}\right)^{\frac{1}{2}n+\lambda} K_{\frac{1}{2}n+\lambda}(m|\vec{z}|),\end{aligned}\quad (3)$$

where K denotes the modified Bessel-functions of second order and Γ is the Gamma function.

III | NEWTON-WIGNER-PRYCE OPERATOR

As already pointed out in the introduction, the appropriate (spatial)-position operator for the Klein-Gordon field is given by the so-called Newton-Wigner-Pryce operator, (Pryce, 1948) and (Newton and Wigner, 1949). For the one-particle case it is given by the following action on a one-particle wave function in momentum space, (Schweber, 1961, Chapter 3c, Equation 35)

$$(X_j \varphi)(\mathbf{p}) = -i \left(\frac{p_j}{2\omega_{\mathbf{p}}^2} + \frac{\partial}{\partial p^j} \right) \varphi(\mathbf{p}), \quad (4)$$

where here we use the covariant representation and normalization, i.e. the observable X_j is represented as a self-adjoint operator w.r.t. the \mathcal{H}_1 -scalar product, i.e. w.r.t. to the measure $d^n \mu(\mathbf{p})$. For the free (relativistic) scalar field the eigenfunctions of the Newton-Wigner-Pryce operator, which are simultaneously the localized wave functions at time $x_0 = 0$, are given by, (Schweber, 1961, Chapter 3, Equation 38)

$$\Psi_{\mathbf{x},0}(\mathbf{p}) = (2\pi)^{-n/2} e^{-i\mathbf{p}\cdot\mathbf{x}} (2\omega_p)^{1/2}.$$

They are of physical importance from the following point of view. Let a particle be in a state $\Phi(\mathbf{p})$ at time $t = 0$, then the probability amplitude that a position measurement will find the particle at \mathbf{x} is given by, (Schweber, 1961, Chapter 3, Equation 44)

$$\langle \bar{\Psi}_{\mathbf{x}}, \Phi \rangle = \int d^n \mu(\mathbf{p}) \Psi_{\mathbf{x},0}(\mathbf{p}) \Phi(\mathbf{p}).$$

In (Sibold and Solard, 2009), (Much, 2013) and (Much, 2015) a second-quantized version of this operator was given. Hence, the k -particle generalization of the Newton-Wigner-Pryce operator can be defined. Particularly, it can be obtained by the Fourier-transformation of the spatial momentum operator.

It is given as a self-adjoint operator on the domain $\bigotimes_{i=1}^k \mathcal{S}(\mathbb{R}^n)$, with $\mathcal{S}(\mathbb{R}^n)$ denoting the Schwartz space, as follows, (Much, 2013; Sibold and Solard, 2009)

$$X_j = -i \int d^n \mathbf{p} a^*(\mathbf{p}) \frac{\partial}{\partial p^j} a(\mathbf{p}), \quad (5)$$

where the second-quantized momentum operator, is given as a self-adjoint operator on the domain

$\bigotimes_{i=1}^k \mathcal{S}(\mathbb{R}^n)$ and in terms of the ladder operators as

$$P_\mu = \int d^n \mathbf{p} p_\mu a^*(\mathbf{p}) a(\mathbf{p})$$

The commutator of the spatial-momentum operator and the NWP-operator is simply the second-quantized Heisenberg-Weyl relation, see (Sibold and Solard, 2009) or (Much, 2015),

$$[X_j, P_k] = -i\eta_{jk}N, \quad (6)$$

where N is the particle-number operator represented in Fock-space as

$$N = \int d^n \mathbf{p} a^*(\mathbf{p}) a(\mathbf{p}). \quad (7)$$

Remark In the last commutator relation the zero component of the coordinate operator is missing. The issue will be addressed in the following section more thoroughly, however let us refer the reader to the following works about this subject and related Ansätze to solve it, (Sibold and Solard, 2009), (Sibold and Burkhard, 2010), (Much, 2015) and (Much and Vergara, 2017).

In what follows we give the form and proof of the coordinate operator in coordinate space.

Lemma 1. *The Newton-Wigner-Pryce operator has the following coordinate space representation,*

$$X_j = \int d^n \mathbf{x} x_j \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x})$$

Demostración. The proof is done by changing from the momentum basis to the coordinate basis, i.e.

$$\begin{aligned} X_j &= -i \int d^n \mathbf{p} a^*(\mathbf{p}) \frac{\partial}{\partial p_j} a(\mathbf{p}) \\ &= -i(2\pi)^{-n} \int d^n \mathbf{p} \int d^n \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \int d^n \mathbf{y} \left(\frac{\partial}{\partial p_j} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\ &= (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} y_j \underbrace{\left(\int d^n \mathbf{p} e^{-ip_k(x-y)^k} \right)}_{(2\pi)^n \delta(\mathbf{x}-\mathbf{y})} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}), \end{aligned}$$

where in the last lines we performed the derivative and integrated over the delta distribution. \square

Next, we want to compare the second-quantized operator with the k -particle extension in literature and draw attention to the distinction between coordinate space and configuration space (see (Schweber, 1961, Chapter 7)). In order to do so let us introduce the configuration space in QFT. For a free scalar field a one particle state is given by

$$|x\rangle = \phi^-(x)|0\rangle = \int d^n \mu(\mathbf{p}) e^{ipx} a_c^*(\mathbf{p})|0\rangle = \int d^n \mu(\mathbf{p}) e^{ipx} |\mathbf{p}\rangle,$$

where $\phi^-(x)$ contains only negative-frequency parts of the Klein-Gordon field at a **space-time** point x . Moreover it should be noted that the following expressions are heuristic, and only make sense when

suitably smeared with a test function. One can think of the configuration space as the space spanned by vectors,

$$|x_1, \dots, x_k\rangle = (k!)^{-1/2} \phi^-(x_1) \cdots \phi^-(x_k) |0\rangle,$$

where the vector $|0\rangle$ is the vacuum. The k -particle Fock-space amplitude for a vector $|\Psi\rangle$ is given by the scalar product with the configurations space vectors, i.e.

$$\Psi^{(k)}(x_1, \dots, x_k) = (k!)^{-1/2} \langle 0 | \phi^+(x_1) \cdots \phi^+(x_k) | \Psi \rangle,$$

where $\phi^+(x)$ contains only positive-frequency parts of the Klein-Gordon field at a point x . From a physical point of view, this is **not** the probability amplitude for finding k -particle at position $\mathbf{x}_1 \cdots \mathbf{x}_k$ at time $x_0 = x_{10} = \cdots x_{k0}$. But it is rather the probability amplitude for finding k -particle at time x_0 . Hence, in order to find the quantity, which gives as the probability amplitude for finding k -particle at positions $\mathbf{x}_1 \cdots \mathbf{x}_k$ at time $x_0 = x_{10} \cdots x_{k0}$, as we did before for one-particle, we introduce the following operator, (Schweber, 1961, Chapter 7, Equation 99),

$$\phi_1(x) = \int d^n \mu(\mathbf{p}) \overline{\Psi_{\mathbf{x}, x_0}(\mathbf{p})} a_c(\mathbf{p}) \quad (8)$$

$$= (2\pi)^{-n/2} \int d^n \mu(\mathbf{p}) e^{ip \cdot x} (2\omega_p)^{1/2} a_c(\mathbf{p}), \quad (9)$$

and apply k of them on $|\Phi^1\rangle$ and the vacuum as follows,

$$\Phi^1((x_0, \mathbf{x}_1), \dots, (x_0, \mathbf{x}_k)) = (k!)^{-1/2} \langle 0 | \phi_1(x_0, \mathbf{x}_1) \cdots \phi_1(x_0, \mathbf{x}_k) | \Phi^1 \rangle.$$

The discussion of amplitudes as probabilities at time x_0 makes sense naturally when the above expressions are smeared, i.e. by integrating over a spatial smearing function at fixed time x_0 . Although we naively introduced in the preliminaries the Fourier transformation of the creation and annihilation operators, we have the following equality

$$\phi_1(x)|_{x_0=0} = \tilde{a}(\mathbf{x}). \quad (10)$$

The equivalence is easily proven by taking the non-covariant normalization and the inverse Fourier-transformation into account. The eigenstates of the second-quantized coordinate operator are created by the action of $\tilde{a}(\mathbf{x}_1), \dots, \tilde{a}(\mathbf{x}_k)$ on the vacuum from the right. Hence, the second-quantized position operator given in Equation (5) agrees with the definition of an operator creating the coordinate space, that is needed for the calculation of probability amplitudes of positions of the particles.

To understand **where** (spatially) the particles are, the coordinate space displays more importance, than the configuration space. Particularly, it is essential to obtain a deeper understanding of QFT's that are defined on non-commutative spaces.

Remark An integral transformation from the configuration space to the coordinate space exists and it is given by,

$$\begin{aligned}
|x\rangle &= \phi^+(x)|0\rangle = \int d^n \mu(\mathbf{p}) e^{ipx} a_c^*(\mathbf{p})|0\rangle \\
&= (2\pi)^{-n/2} \int d^n \mathbf{y} \left(\int d^n \mu(\mathbf{p}) \sqrt{2\omega_{\mathbf{p}}} e^{i(\omega_{\mathbf{p}}x^0 + p_k(x-y)^k)} \right) \tilde{a}^*(\mathbf{y})|0\rangle,
\end{aligned}$$

where in the last line the Fourier-transformation was used.

IV | POINCARÉ GROUP

In this section, we take the representations of the Poincaré group in terms of QFT Fock-space operators and perform a base change to the coordinate space.

Lemma 2. *The zero component, i.e. the operator generating time-translations, has the following representation in coordinate space,*

$$P_0 = \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) (\tilde{\omega} * \tilde{a})(\mathbf{x}),$$

where $*$ denotes the convolution and the function $\tilde{\omega}(\mathbf{x})$ is defined as

$$\tilde{\omega}(\mathbf{x}) := -2(2\pi)^{-\frac{n+1}{2}} \left(\frac{m}{|\mathbf{x}|} \right)^{\frac{n+1}{2}} K_{\frac{n+1}{2}}(m|\mathbf{x}|).$$

The momentum operator, that is responsible for spatial translations takes in coordinate space the following form,

$$P_j = i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) \frac{\partial}{\partial x^j} \tilde{a}(\mathbf{x}).$$

Demostración. Let us first prove the base change for the spatial part of the momentum operator. The base change is straight forward and it is done as in the proof for the coordinate operator (see Lemma 4),

$$\begin{aligned}
P_j &= \int d^n \mathbf{p} p_j a^*(\mathbf{p}) a(\mathbf{p}) \\
&= (2\pi)^{-n} \int d^n \mathbf{p} p_j \int d^n \mathbf{x} e^{-ip_l x^l} \tilde{a}^*(\mathbf{x}) \int d^n \mathbf{y} e^{ip_k y^k} \tilde{a}(\mathbf{y}) \\
&= (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} \left(\int d^n \mathbf{p} p_j e^{-ip_k(x-y)^k} \right) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\
&= -(2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} i \frac{\partial}{\partial y^j} \left(\int d^n \mathbf{p} e^{-ip_k(x-y)^k} \right) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\
&= i \iint d^n \mathbf{x} d^n \mathbf{y} \delta^n(\mathbf{x} - \mathbf{y}) \tilde{a}^*(\mathbf{x}) \frac{\partial}{\partial y^j} \tilde{a}(\mathbf{y}) \\
&= i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) \frac{\partial}{\partial x^j} \tilde{a}(\mathbf{x}),
\end{aligned}$$

where in the last lines we performed a partial integration and integrated over the delta function. Next, we transform the zero component into the coordinate space. This transformation requires more work and in particular we use Formula (3),

$$\begin{aligned}
P_0 &= \int d^n \mathbf{p} \omega_{\mathbf{p}} a^*(\mathbf{p}) a(\mathbf{p}) \\
&= (2\pi)^{-n} \int d^n \mathbf{p} \sqrt{|\mathbf{p}|^2 + m^2} \int d^n \mathbf{x} e^{-i p_l x^l} \tilde{a}^*(\mathbf{x}) \int d^n \mathbf{y} e^{i p_k y^k} \tilde{a}(\mathbf{y}) \\
&= (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} \left(\int d^n \mathbf{p} \sqrt{|\mathbf{p}|^2 + m^2} e^{-i p_k (x-y)^k} \right) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\
&= -2(2\pi)^{-\frac{n+1}{2}} \iint d^n \mathbf{x} d^n \mathbf{y} \left(\frac{m}{|\mathbf{x} - \mathbf{y}|} \right)^{\frac{n+1}{2}} K_{\frac{n+1}{2}}(m|\mathbf{x} - \mathbf{y}|) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}),
\end{aligned}$$

where the Fourier transformation in the last step can be found in (Gel'fand and Shilov, 1964, Chapter III, Section 2.8). \square

An interesting object that appeared naturally in (Much, 2015) is that of the velocity operator. It can be obtained in two different ways. The first path is guided by intuition. Since the operator must be the second quantization of the spatial velocity for a relativistic particle we have the following expression,

$$V_j = \int d^n \mathbf{p} \frac{P_j}{\omega_{\mathbf{p}}} a^*(\mathbf{p}) a(\mathbf{p})$$

The second more profound approach is given by the Heisenberg equation of motion,

$$[P_0, X_j] = -iV_j. \quad (11)$$

The results of the two paths agree and it was proven in (Much, 2015). Since we have expressions for the momentum and coordinate operator in the spatial space, we can express the velocity operator in the coordinate space by taking the commutator (11). Hence, the second path is from a calculative point of view easier and the explicit result is given in the following lemma.

Lemma 3. *The velocity operator expressed in terms of ladder operators of the coordinate space is given by,*

$$V_j = -i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) (\tilde{\omega}_j * \tilde{a})(\mathbf{x}), \quad (12)$$

with vector-valued function $\tilde{\omega}_j$ defined as $\tilde{\omega}_j(\mathbf{x}) := 2(2\pi)^{-\frac{n+1}{2}} \left(\frac{m}{|\mathbf{x}|} \right)^{\frac{n+1}{2}} K_{\frac{n+1}{2}}(m|\mathbf{x}|) x_j$ and where $*$ denotes the convolution.

Demostración. The proof is straight-forward and it is done by explicitly calculating the Heisenberg equation of motion in coordinate space, i.e.

$$\begin{aligned}
[P_0, X_j] &= \iint d^n \mathbf{x} d^n \mathbf{z} z_j [\tilde{a}^*(\mathbf{x}) (\tilde{\omega}_j * \tilde{a})(\mathbf{x}), \tilde{a}^*(\mathbf{z}) \tilde{a}(\mathbf{z})] \\
&= \iint d^n \mathbf{x} d^n \mathbf{y} d^n \mathbf{z} z_j \tilde{\omega}(x-y) \underbrace{[\tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}), \tilde{a}^*(\mathbf{z}) \tilde{a}(\mathbf{z})]}_{-\delta(\mathbf{x}-\mathbf{z}) \tilde{a}^*(\mathbf{z}) \tilde{a}(\mathbf{y}) + \delta(\mathbf{y}-\mathbf{z}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{z})} \\
&= - \iint d^n \mathbf{x} d^n \mathbf{y} (x-y)_j \tilde{\omega}(x-y) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}).
\end{aligned}$$

□

The velocity operator expressed in the coordinate space is useful for the boost operators. In particular, the boosts of the restricted Lorentz group have the velocity operator explicitly in their respective representations.

Next, we turn to the generators of the proper orthochronous Lorentz group \mathcal{L}_+^\uparrow , consisting of boosts and rotations which are given in the momentum space as, (Itzykson and Zuber, 1980, Equation 3.54), (Sibold and Solard, 2009)

$$M_{j0} = i \int d^n \mathbf{p} a^*(\mathbf{p}) \left(\frac{p_j}{2\omega_{\mathbf{p}}} - \omega_{\mathbf{p}} \frac{\partial}{\partial p^j} \right) a(\mathbf{p}), \quad (13)$$

$$M_{ik} = i \int d^n \mathbf{p} a^*(\mathbf{p}) \left(p_i \frac{\partial}{\partial p^k} - p_k \frac{\partial}{\partial p^i} \right) a(\mathbf{p}). \quad (14)$$

The next theorem displays the importance of the NWP-operator. In particular, the generators of boost and rotations can be represented by second quantized (denoted by $d\Gamma(\cdot)$ see (Reed and Simon, 1975, Chapter X.7)) products of the momentum and NWP-operator.

Theorem 4. *For the massive scalar field, generators of the proper orthochronous Lorentz-group \mathcal{L}_+^\uparrow can be represented in terms of products of the NWP-operator and the relativistic momentum operator as follows,*

$$M_{0j} = \frac{1}{2} (d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)), \quad M_{ik} = d\Gamma(X_i P_k) - d\Gamma(X_k P_i). \quad (15)$$

Demostración. We start the proof for boosts where we have,

$$\begin{aligned} M_{0j} &= \frac{1}{2} (d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)) = \frac{1}{2} d\Gamma([X_j, P_0]) + d\Gamma(P_0 X_j) \\ &= \frac{i}{2} d\Gamma(V_j) + d\Gamma(P_0 X_j), \end{aligned}$$

where in the last lines we used the Equation of motion given by Formula (11). The proof for rotations is obvious and hence it is omitted. □

The former Theorem conveys the fact that we can define the generator of the Lorentz-group by using the coordinate operator and the momentum operator. Therefore, the canonical commutation relations with addition of the zero component of momentum can be used to define the group of relativity. This fact, is, in our opinion, an additional argument for the physical sense of the NWP-operator.

Moreover, the representation of Lorentz generators by using the NWP and momentum operator induces more transparency into the non-covariant behavior of the NWP-operator with regards to boosts. The following calculations clarify the former statement,

$$\begin{aligned} [M_{0j}, P_0] &= \frac{1}{2} [(d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)), d\Gamma(P_0)] \\ &= \frac{1}{2} (d\Gamma([X_j, P_0] P_0) + d\Gamma(P_0 [X_j, P_0])) \\ &= i d\Gamma(V_j P_0) = i d\Gamma(P_j) = iP_j, \end{aligned}$$

where in the last lines we used $[d\Gamma(A), d\Gamma(B)] = d\Gamma([A, B])$ and the Heisenberg-equation (see Equation 11). Moreover, we used the representation of the velocity operator as $V_j = d\Gamma(P_0^{-1}P_j)$. Next, the commutator of boosts with the spatial momentum is calculated,

$$\begin{aligned} [M_{0j}, P_k] &= \frac{1}{2} [(d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)), d\Gamma(P_k)] \\ &= \frac{1}{2} (d\Gamma([X_j, P_k] P_0) + d\Gamma(P_0 [X_j, P_k])) \\ &= -i\eta_{jk} P_0, \end{aligned}$$

where in the last lines we used the second-quantization of the unit operator to be the particle number operator, i.e. $d\Gamma(\mathbb{1}_{\mathcal{H}}) = N$ and the explicit canonical commutation relations, see Equation (6). Next, we calculate the commutator which essentially answers the question of covariance w.r.t. the NWP-operator,

$$\begin{aligned} [M_{0j}, X_k] &= \frac{1}{2} [(d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)), d\Gamma(X_k)] \\ &= \frac{1}{2} (d\Gamma(X_j [P_0, X_k]) + d\Gamma([P_0, X_k] X_j)) \\ &= \frac{i}{2} (d\Gamma(X_j V_k) + d\Gamma(V_k X_j)). \end{aligned}$$

The former commutator has no resemblance to a covariant commutator-like object as $[M_{0j}, X_k] = -i\eta_{jk} X_0$. However, the boost operator can be translated in time by using the generators of the translation group,

$$\begin{aligned} e^{ix^0 P_0} M_{j0} e^{-ix^0 P_0} &= M_{j0} + ix^0 [P_0, M_{j0}] + \frac{i^2}{2!} (x^0)^2 \underbrace{[P_0, [P_0, M_{j0}]]}_{=0} + \dots \\ &= M_{j0} + x^0 P_j, \end{aligned}$$

with $x^0 \in \mathbb{R}$ and where the Baker-Campbell-Hausdorff formula and the explicit Poincaré algebra was used. Therefore, the time-dependent boost-operator can be represented as a symmetric product of the second quantized coordinate, the time and the momentum operators,

$$\begin{aligned} M_{0j} &= \frac{1}{2} (d\Gamma(X_j P_0) + d\Gamma(P_0 X_j) + x_0 d\Gamma(P_0) + d\Gamma(P_0) x_0) \\ &= \frac{1}{2} (d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)) + x_0 d\Gamma(P_0). \end{aligned}$$

By using the time-translated boost operator we obtain for the commutator with the NWP-operator ,

$$\begin{aligned} [M_{j0} + x^0 P_j, X_k] &= \frac{i}{2} (d\Gamma(X_j V_k) + d\Gamma(V_k X_j)) + x^0 [P_j, X_k] \\ &= i\eta_{jk} x^0 N + \frac{i}{2} (d\Gamma(X_j V_k) + d\Gamma(V_k X_j)), \end{aligned}$$

which has a resemblance to a covariant operator. From the view point of representing the Lorentz generators by the use of the NWP- and the momentum operator, the non-covariance issue becomes clear on a more profound level. Commutator relations for the rotations are obvious and hence we shall

omit them. However, from the representation of these operators, it becomes obvious why both the NWP and the momentum operator transform covariantly w.r.t rotations. In particular, generators of rotations have the well-known complementarity between momentum and position.

Lemma 5. *Generators of the proper orthochronous Lorentz group \mathcal{L}_+^\uparrow expressed in the terms of ladder operators of the free massive scalar field are represented in the coordinate space as,*

$$M_{j0} = \frac{1}{2} \iint d^n \mathbf{x} d^n \mathbf{y} (x+y)_j \tilde{\omega}(\mathbf{x}-\mathbf{y}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y})$$

The operator of rotations takes the familiar form in the coordinate space,

$$M_{ik} = i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) \left(x_i \frac{\partial}{\partial x^k} - x_k \frac{\partial}{\partial x^i} \right) \tilde{a}(\mathbf{x}).$$

Demostración. We start by calculating the base change of the rotation generators of the Lorentz group \mathcal{L}_+^\uparrow ,

$$\begin{aligned} M_{ik} &= i \int d^n \mathbf{p} a^*(\mathbf{p}) \left(p_i \frac{\partial}{\partial p^k} - p_k \frac{\partial}{\partial p^i} \right) a(\mathbf{p}) \\ &= i(2\pi)^{-n} \int d^n \mathbf{p} \int d^n \mathbf{x} e^{-ip_l x^l} \tilde{a}^*(\mathbf{x}) \left(p_i \frac{\partial}{\partial p^k} - p_k \frac{\partial}{\partial p^i} \right) \int d^n \mathbf{y} e^{ip_r y^r} \tilde{a}(\mathbf{y}) \\ &= -i(2\pi)^{-n} \int d^n \mathbf{p} \int d^n \mathbf{x} e^{-ip_l x^l} \tilde{a}^*(\mathbf{x}) \int d^n \mathbf{y} \left(y_i \frac{\partial}{\partial y^k} - y_k \frac{\partial}{\partial y^i} \right) e^{ip_r y^r} \tilde{a}(\mathbf{y}) \\ &= i(2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} \left(\int d^n \mathbf{p} e^{-ip_l(x-y)^l} \right) \tilde{a}^*(\mathbf{x}) \left(y_i \frac{\partial}{\partial y^k} - y_k \frac{\partial}{\partial y^i} \right) \tilde{a}(\mathbf{y}) \\ &= i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) \left(x_i \frac{\partial}{\partial x^k} - x_k \frac{\partial}{\partial x^i} \right) \tilde{a}(\mathbf{x}), \end{aligned}$$

where in the last lines we performed a partial integration and integrated over a delta function. Next, we turn to the Lorentz boosts. The first term is simply the velocity operator times $\frac{i}{2}$. Hence, we focus here only on the second part,

$$\begin{aligned} &-i \int d^n \mathbf{p} a^*(\mathbf{p}) \omega_{\mathbf{p}} \frac{\partial}{\partial p^j} a(\mathbf{p}) \\ &= -i \int d^n \mathbf{p} \int d^n \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \omega_{\mathbf{p}} \frac{\partial}{\partial p^j} \int d^n \mathbf{y} e^{ip_k y^k} \tilde{a}(\mathbf{y}) \\ &= (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} \left(\int d^n \mathbf{p} \sqrt{|\mathbf{p}|^2 + m^2} e^{-ip_k(x-y)^k} \right) y_j \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\ &= -2(2\pi)^{-\frac{n+1}{2}} \iint d^n \mathbf{x} d^n \mathbf{y} y_j \left(\frac{m}{|\mathbf{x}-\mathbf{y}|} \right)^{\frac{n+1}{2}} K_{\frac{n+1}{2}}(m|\mathbf{x}-\mathbf{y}|) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\ &= \iint d^n \mathbf{x} d^n \mathbf{y} y_j \tilde{\omega}(\mathbf{x}-\mathbf{y}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}), \end{aligned}$$

where in the last lines we performed the differentiation and used Integral (3). □

Now the result for the rotations is not surprising and is well-known from quantum mechanics, i.e. for one-particle. The interesting and unknown result is that of the Lorentz-boosts. It is a representation of the boost operator in terms of the spatial coordinate space. From the form of the boost generator it also becomes more clear why a covariant coordinate operator, i.e. a zero component (besides issues with Pauli's theorem) X_0 cannot exist. This can be seen by taking the commutator of the Lorentz boost and the coordinate operator,

$$\begin{aligned} [M_{j0}, X_k] &= \frac{1}{2} \iiint d^n \mathbf{x} d^n \mathbf{y} d^n \mathbf{z} (x+y)_j z_k \tilde{\omega}(\mathbf{x}-\mathbf{y}) [\tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}), \tilde{a}^*(\mathbf{z}) \tilde{a}(\mathbf{z})] \\ &= -\frac{1}{2} \iiint d^n \mathbf{x} d^n \mathbf{y} (x+y)_j (x-y)_k \tilde{\omega}(\mathbf{x}-\mathbf{y}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\ &\neq -i \delta_{jk} X_0. \end{aligned}$$

Moreover, the complementarity in an operational sense, i.e. by replacing multiplication operators with derivatives and vice versa, is broken by the boost operators. This can be easily seen due to their explicit form in the coordinate space.

In Section III we gave the explicit operator (see Equation (8)) to calculate the probability amplitude for finding k -particle at positions $\mathbf{x}_1 \cdots \mathbf{x}_k$ at time $x_0 = x_{10} = \cdots x_{k0}$. By setting the time component $x_0 = 0$ equal to zero we were able to match this expression with our ladder operator in coordinate space. Hence, we could (or should) have calculated the more general and explicit expressions for the operators by keeping the time component. To resolve this issue, time-translations can be performed by using the unitary adjoint action of the zero component of the momentum. This fact is composed as a result in the form of a proposition.

Proposition 6. *The operator ϕ_1 given as,*

$$\phi_1(x) = (2\pi)^{-n/2} \int d^n \mathbf{p} e^{ip \cdot x} a(\mathbf{p}),$$

can be expressed by using time-translations of the Fourier-transformed annihilation operator \tilde{a} as follows

$$\phi_1(x) = \phi_1(x^0, \mathbf{x}) = e^{ix^0 P_0} \tilde{a}(\mathbf{x}) e^{-ix^0 P_0}, \quad x^0 \in \mathbb{R}.$$

Demostración. The proof is performed by using the inverse Fourier-transformation of \tilde{a} , i.e.

$$\begin{aligned} e^{ix^0 P_0} \tilde{a}(\mathbf{x}) e^{-ix^0 P_0} &= (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-ip_k x^k} e^{ix^0 P_0} a(\mathbf{p}) e^{-iy^0 P_0} \\ &= (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-ip_k x^k} e^{-ip_0 x^0} a(\mathbf{p}), \end{aligned}$$

where in the last lines we used the adjoint action of the translation group on the annihilation operator, see (Schweber, 1961, Chapter 7, Equation 61). This agrees with the expression given in Equation (8). The explicit expression of this operator in terms of the coordinate space ladder operators is given in the next section. \square

Hence, instead of using the time-independent transformations using the ladder operators in coordinate space, we could have used the product of $\phi_1(x^0, \mathbf{x})$ and $\phi_1(x^0, \mathbf{y})$ or we can simply take our obtained

expressions and perform a time-translation. However, from the Baker-Campbell-Hausdorff formula and the particular algebra of the Poincaré group it follows that the momentum operator and the spatial rotations are invariant under time-translations. Hence, the only term that changes under time-translation is the boost operator and this is easily calculated,

$$e^{ix^0 P_0} M_j e^{-ix^0 P_0} = M_j + x^0 P_j,$$

using the Baker-Campbell-Hausdorff formula and the Poincaré algebra.

Although, neither the coordinate operator nor the respective eigenstates are Lorentz-covariant, the Poincaré operators that are translated in this work obey the covariant transformation property. In particular, the Lorentz-covariance does not depend on the specific base that is chosen. This statement is proven by re-transforming the ladder operators from coordinate space into momentum space, and taking the appropriate transformation into account. This is obvious and it simply follows from the fact that the operators are covariant, invariant of the representation they are given in. Hence, even though we use non-covariant eigenstates, generators of the Poincaré group written in coordinate space respect the relativistic covariance.

V | POINCARÉ TRANSFORMATIONS OF THE COORDINATE SPACE

In this section we investigate Poincaré transformations, for the massive case, of the Fourier-transformed ladder operators. In particular, we calculate how the operator ϕ_1 , which is the object used to calculate probability amplitudes, transforms under space-time translations and pure rotations. The motivation herein is to give the proper transformation behavior of probability amplitudes under changes of frame. In particular, given two frames related by space-time translations or pure rotations, both must be able to calculate and compare the probability amplitude w.r.t. the frame change. We purposely neglect boosts since the spatial coordinate space is not Lorentz-covariant and hence the question of how boosts act is not well-defined from the very beginning.

In the forthcoming calculations, we work in the physical relevant dimension $d = 4$. Moreover, we define the unitary operator that generates transformations of the orthochronous proper Poincaré group, $\mathcal{P}_+^\uparrow = \mathcal{L}_+^\uparrow \times \mathbb{R}^4$ as $U(y, \Lambda)$ transforming the creation and annihilation operators as, (Schweber, 1961, Chapter 7),

$$U(y, \Lambda) a(\mathbf{p}) U(y, \Lambda)^{-1} = \sqrt{\frac{\Omega_{\Lambda \mathbf{p}}}{\omega_{\mathbf{p}}}} e^{-i(\Lambda p)_\mu y^\mu} a(\Lambda \mathbf{p}), \quad (y, \Lambda) \in \mathcal{P}_+^\uparrow, \quad (16)$$

$$U(y, \Lambda) a^*(\mathbf{p}) U(y, \Lambda)^{-1} = \sqrt{\frac{\Omega_{\Lambda \mathbf{p}}}{\omega_{\mathbf{p}}}} e^{i(\Lambda p)_\mu y^\mu} a^*(\Lambda \mathbf{p}), \quad (y, \Lambda) \in \mathcal{P}_+^\uparrow \quad (17)$$

To simplify calculations with regard to transformations of the operator ϕ_1 , we first give the adjoint actions on the Fourier-transformed operators and afterwards perform a time-translation, see Proposition 6. By using the former results, we calculate transformations in the momentum space and perform an inverse Fourier-transformation in order to get the respective coordinate space transformations.

Lemma 7. *The Fourier-transformed annihilation operator \tilde{a} transforms under the adjoint action of the translation group as follows,*

$$U(y, \mathbb{I})\tilde{a}(\mathbf{x})U(y, \mathbb{I})^{-1} = \frac{iy^0}{2\pi^2} \int d^3\mathbf{z} \left(\frac{m^2 K_2(m\sqrt{|\mathbf{x} + \mathbf{y} - \mathbf{z}|^2 - (y^0)^2})}{|\mathbf{x} + \mathbf{y} - \mathbf{z}|^2 - (y^0)^2} \right) \tilde{a}(\mathbf{z}),$$

where $y \in \mathbb{R}^4$ and $*$ denotes the convolution.

Demostración. Since, the momentum operators commute we can first calculate the spatial transformations, which are easier, and then the time-translations, i.e.

$$U(y, \mathbb{I})\tilde{a}(\mathbf{x})U(y, \mathbb{I})^{-1} = U(y^0, \mathbb{I})U(\mathbf{y}, \mathbb{I})\tilde{a}(\mathbf{x})U(\mathbf{y}, \mathbb{I})^{-1}U(y^0, \mathbb{I})^{-1}.$$

Therefore, we first investigate the following expression,

$$\begin{aligned} U(\mathbf{y}, \mathbb{I})\tilde{a}(\mathbf{x})U(\mathbf{y}, \mathbb{I})^{-1} &= e^{iy^k P_k} \tilde{a}(\mathbf{x}) e^{-iy^k P_k} \\ &= (2\pi)^{3/2} \int d^3\mathbf{p} e^{-ip_k x^k} U(\mathbf{y}, \mathbb{I}) a(\mathbf{p}) U(\mathbf{y}, \mathbb{I})^{-1} \\ &= (2\pi)^{-3/2} \int d^3\mathbf{p} e^{-ip_k (x+y)^k} a(\mathbf{p}) = \tilde{a}(\mathbf{x} + \mathbf{y}), \end{aligned}$$

where in the last lines we used the inverse Fourier-transformation (see Equation 2) and the well-known action of translations in the momentum space, see (Schweber, 1961, Chapter 7, Equation 61). Next, we calculate the action of time-translations, which is given by the following expression,

$$\begin{aligned} U(y^0, \mathbb{I})\tilde{a}(\mathbf{x})U(y^0, \mathbb{I})^{-1} &= e^{iy^0 P_0} \tilde{a}(\mathbf{x}) e^{-iy^0 P_0} \\ &= (2\pi)^{-3/2} \int d^3\mathbf{p} e^{-ip_k x^k} e^{-iy^0 \omega_{\mathbf{p}}} a(\mathbf{p}) \\ &= (2\pi)^{-3} \int d^3\mathbf{z} \left(\int d^3\mathbf{p} e^{-ip_k (x-z)^k} e^{-i\omega_{\mathbf{p}} y^0} \right) \tilde{a}(\mathbf{z}) \\ &= \frac{iy^0}{2\pi^2} \int d^3\mathbf{z} \left(\frac{m^2 K_2(m\sqrt{|\mathbf{x} - \mathbf{z}|^2 - (y^0)^2})}{|\mathbf{x} - \mathbf{z}|^2 - (y^0)^2} \right) \tilde{a}(\mathbf{z}), \end{aligned}$$

where in the former equation we used (Peskin and Schroeder, 1995, Chapter 2) and (Gradshteyn et al., 2007, Equation 3.914) to solve the integral. Equivalent considerations can be done for the Fourier transformed creation operator \tilde{a}^* . By using both transformations the proof is completed. \square

The translation acting on the coordinate space in spatial direction was as expected a translation in the spatial space as well. However, the difference in the spatial space becomes visible when we perform a translation in time. Since, there is no time component the time-translation transforms, and therefore acts, on the spatial space in a more complex manner. It is interesting since this is a physical important expression. Essentially, it tells us how the spatial coordinate space (which is made of k -ladder operators, see Equation 10), transforms under a time-translation. Moreover, with regards to the time-translated coordinate annihilation operator \tilde{a} , which corresponds to ϕ_1 , an explicit expression in coordinate space was given. The next object of interest is the Lorentz-transformation of the Fourier-transformed ladder operators.

Lemma 8. Let Λ_R denote the matrices of the Lorentz group which represent pure rotations. They are given by

$$\Lambda_R = \begin{pmatrix} 1 & 0 \\ 0 & R \end{pmatrix}, \quad R \in SO(3),$$

where $SO(3)$ is the group of real, orthogonal, 3×3 matrices with determinant one. Then, the Fourier-transformed ladder operator \tilde{a} transforms under pure rotations as follows

$$U(0, \Lambda_R) \tilde{a}(\mathbf{x}) U(0, \Lambda_R)^{-1} = \tilde{a}(R\mathbf{x}).$$

Demostración. The action of pure spatial rotations on the the Fourier-transformed annihilation operator is calculated by using Equation (17),

$$\begin{aligned} U(0, \Lambda_R) \tilde{a}(\mathbf{x}) U(0, \Lambda_R)^{-1} &= (2\pi)^{-3/2} \int d^3 \mathbf{p} e^{-i p_k x^k} U(0, \Lambda_R) a(\mathbf{p}) U(0, \Lambda_R)^{-1} \\ &= (2\pi)^{-3/2} \int d^3 \mathbf{p} e^{-i p_k x^k} \sqrt{\frac{\omega_{R\mathbf{p}}}{\omega_{\mathbf{p}}}} a(R\mathbf{p}) \\ &= (2\pi)^{-3/2} \int d^3 \mathbf{p} e^{-i p_k (R\mathbf{x})^k} a(\mathbf{p}), \end{aligned}$$

in the last lines we used the transformation behavior of the non-covariant momentum ladder operators (see (Schweber, 1961, Chapter 7, Equation 57)) and the orthogonality of R for pure spatial rotations. The proof for the Fourier-transformed creation operator \tilde{a}^* can be done equivalently as for the annihilation operator. However, the transformations for \tilde{a}^* can be as well obtained by taking the adjoint of the Fourier-transformed annihilation operator. \square

Next, by using the former lemmas we calculate the transformational behavior of $\phi_1(x)$ under the whole group of translations and pure rotations.

Theorem 9. The operator $\phi_1(x)$ transforms in a covariant manner under space-time translations and pure rotations, i.e.

$$U(y, \Lambda_R) \phi_1(x) U(y, \Lambda_R)^{-1} = \phi_1(x_0 + y_0, R\mathbf{x} + \mathbf{y}), \quad (y, \Lambda_R) \in \mathcal{P}_+^\uparrow.$$

Moreover, the explicit transformation of the Fourier-transformed operator \tilde{a} under the action of the subgroup $(y, \Lambda_R) \in \mathcal{P}_+^\uparrow$ is in coordinate space given as,

$$U(y, \Lambda_R) \tilde{a}(\mathbf{x}) U(y, \Lambda_R)^{-1} = \frac{iy^0}{2\pi^2} \left(\frac{m^2 K_2(m\sqrt{|\mathbf{x} + \mathbf{y}|^2 - (y^0)^2})}{|\mathbf{x} + \mathbf{y}|^2 - (y^0)^2} \right) * \tilde{a}(\mathbf{x} + \mathbf{y}).$$

Demostración. By using Proposition 6 we can write the operator $\phi_1(x)$ in terms of the Fourier-transformed operator, i.e.

$$U(y, \Lambda_R) \phi_1(x) U(y, \Lambda_R)^{-1} = U(y, \Lambda_R) U(x^0, \mathbb{I}) \tilde{a}(\mathbf{x}) U(x^0, \mathbb{I})^{-1} U(y, \Lambda_R)^{-1},$$

since time-translations commute with space-time translations and pure rotations and since we have $U(y, \Lambda_R) = U(y, \mathbb{I}) U(0, \Lambda_R)$, they allow us to rewrite the former expression as

$$\begin{aligned}
& U(x^0, \mathbb{I}) U(y, \Lambda_R) \tilde{a}(\mathbf{x}) U(y, \Lambda_R)^{-1} U(x^0, \mathbb{I})^{-1} \\
&= U(x^0, \mathbb{I}) U(y, \mathbb{I}) U(0, \Lambda_R) \tilde{a}(\mathbf{x}) U(0, \Lambda_R)^{-1} U(y, \mathbb{I})^{-1} U(x^0, \mathbb{I})^{-1} \\
&= U(x^0, \mathbb{I}) U(y, \mathbb{I}) \tilde{a}(R\mathbf{x}) U(y, \mathbb{I})^{-1} U(x^0, \mathbb{I})^{-1} \\
&= U(x^0 + y^0, \mathbb{I}) \tilde{a}(R\mathbf{x} + \mathbf{y}) U(x^0 + y^0, \mathbb{I})^{-1} \\
&= \phi_1(x_0 + y_0, R\mathbf{x} + \mathbf{y}),
\end{aligned}$$

where in the last lines we used Proposition 6 and the Lemmas 7 and 8, which give the transformational behavior of the coordinate space operator \tilde{a} . \square

VI | CONCLUSION AND OUTLOOK

In this paper we performed a base change of the Poincaré group for a massive scalar field, into the coordinate space. This was done by expressing the creation and annihilation operators, via Fourier-transformation, in terms of the coordinate basis.

One particular interesting, but expected, aspect of our results was the representation of the spatial momentum and the Lorentzian infinitesimal generators of rotations. The one-particle spatial momentum operator and the rotation operators had the same form as in standard quantum mechanics. This is due to the fact that the energy condition, i.e. the choice of $\omega_{\mathbf{p}}$, does not explicitly enter the expressions for these particular observables. Nevertheless, for time-translations and boosts we notice the difference and in particular the complementarity between the \mathbf{x} -space and the momentum space is broken.

Concerning the specific actions of the Poincaré group on this space we obtained important results. The action of the spatial translations and rotations on the coordinate space were as expected. However, the interesting result occurred when the time-translation was involved. In a sense, these transformations are a way to fully understand what happens in the spatial coordinate space, when transformations involving time come into play. The restriction to this particular space is a by-product of the mass shell condition that restricted a d -dimensional momentum space to a $(d - 1)$ -dimensional subspace. Hence, by using the transformations in coordinate space one can calculate the probability of finding particles in certain spatial-positions at time x_0 after an explicit time-translation was performed.

In this work, we translated the boost operators into the coordinate space. Moreover, the adjoint action of the Fourier-transformed operators w.r.t. the boosts was not calculated. Generally, such a calculation is not well-posed since boosts mix time and space in a non-trivial manner. And as hence, the eigenstates of the Newton-Wigner operator are not Lorentz-covariant the calculation of a boost on these particular spaces is from a physical point of view considered problematic.

We performed all base changes by using the massive scalar field. However, a possible extension to this framework can be done with regard to other fields, as for example considering the conformal group expressed in terms of the massless scalar field. This is work in progress.

I | REFERENCIAS

Bratteli, O. and Robinson, D. (1996). Operator algebras and quantum statistical mechanics. Vol. 2: Equilibrium states. Models in quantum statistical mechanics.

- Gel'fand, I. and Shilov, G. (1964). *Generalized functions*. Generalized Functions. Academic Press.
- Gradshteyn, I. S., Ryzhik, I. M., Jeffrey, A., Zwillinger, D., and Scripta Technica, I. (2007). *Table of integrals, series, and products*. Elsevier, Amsterdam, Boston, Paris, et al.
- Itzykson, C. and Zuber, J. (1980). Quantum field theory.
- Jordan, T. F. (1980). Simple derivation of the Newton-Wigner position operator. *J. Math. Phys.*, 21:2028.
- Much, A. (2013). Quantum Spacetime from QM and QFT. *PhD thesis*.
- Much, A. (2015). Relativistic Corrections to the Moyal-Weyl Spacetime. *Journal of Mathematical Physics*, 56(2):022301.
- Much, A. and Vergara, J. D. (2017). A Poincaré Covariant Noncommutative Spacetime?
- Newton, T. D. and Wigner, E. P. (1949). Localized states for elementary systems. *Rev. Mod. Phys.*, 21:400–406.
- Peskin, M. E. and Schroeder, D. V. (1995). *An introduction to quantum field theory*. Advanced book program. Westview Press Reading (Mass.), Boulder (Colo.). Autre tirage : 1997.
- Pryce, M. H. L. (1948). The mass-centre in the restricted theory of relativity and its connexion with the quantum theory of elementary particles. *Proc. Royal Soc. London, Ser. A*, 195:62.
- Reed, M. and Simon, B. (1975). *Methods of Modern Mathematical Physics. 2. Fourier Analysis, Selfadjointness*. Gulf Professional Publishing.
- Schweber, S. S. (1961). An introduction to relativistic quantum field theory.
- Sibold, K. and Burkhard, E. (2010). Conformal algebra on Fock space and conjugate pairs of operators. *Phys.Rev.*, D82:104012.
- Sibold, K. and Solard, G. (2009). Conjugate variables in quantum field theory: The basic case. *Phys.Rev.*, D80:124041.