

TCC No Masiva y el Operador Newton-Wigner

Massless QFT and the Newton-Wigner Operator

A. MUCH¹

Recibido: 2 de febrero de 2018/ Aceptado: 17 de abril de 2018

¹Centro de Ciencias Matemáticas, Universidad Nacional Autónoma de México, C.P. 58190, Morelia, Michoacán México, email: amuch@matmor.unam.mx

Agradecimientos

El autor está en deuda con el Prof. K. Sibold por dar comienzo a las preguntas conceptuales profundas de este trabajo. Además, quisieramos agradecer al Prof. K. Sibold y S. Pottel por muchas discusiones fructíferas durante las diferentes etapas de este artículo. En lo que respecta a los dominios de las transformaciones de Lorentz estoy en deuda con el Prof. Norbert Dragon. Las correcciones hechas por el Dr. Z. Much son reconocidas agradecidamente.

Acknowledgment

The author is indebted to Prof. K. Sibold for initiating deep conceptual questions of this work. Furthermore, we would like to thank Prof. K. Sibold and S. Pottel for many fruitful discussions during different stages of this paper. With regards to the domains of the Lorentz-boosts am indebted to Prof. Norbert Dragon. The corrections of Dr. Z. Much are thankfully acknowledged.

In this work, the second-quantized version of the spatial-coordinate operator, known as the Newton-Wigner-Pryce operator, is explicitly given w.r.t. the massless scalar field. Moreover, transformations of the conformal group are calculated on eigenfunctions of this operator in order to investigate the covariance group w.r.t. probability amplitudes of localizing particles.

En este trabajo, se da explícitamente la versión de segunda cuantización del operador de coordenadas espaciales, conocido como el operador de Newton-Wigner-Pryce, con referencia al campo escalar no masivo. Además las transformaciones del grupo conforme se calculan sobre las eigenfunciones de este operador a fin de investigar el grupo de covariancia con referencia a las amplitudes de probabilidad de partículas en localización.

PALABRAS CLAVES

Fundamentos de la Teoría Cuántica de Campos, Segunda Cuantización, Operador de Coordenadas, Operador Newton-Wigner, TCC en el espacio de coordenadas, Transformación de Fourier.

KEYWORDS

Foundations of Quantum Field Theory, Second Quantization, Coordinate Operator, Newton-Wigner Operator, QFT in coordinate space, Fourier-transformation

PACS

11.10.Cd, 11.10.Nx, 11.30.Cp, 11.90.+t, 03.65.-w, 03.65.Fd, 03.70.+k, 03.30.+p

I | INTRODUCTION

Localization is from a mathematical and physical point of view considered to be an unresolved issue in a relativistic context, see for example (16), (6), (9), (10) and references therein. The concept is well-defined from a non-relativistic quantum-mechanical (QM) point of view. However, by combining concepts of QM and relativity, from which the frame-work of quantum field theory (QFT) emerged, the localization issue was never completely resolved. This is not merely an issue of construction but points

* Esta obra está bajo una licencia Creative Commons Reconocimiento - NoComercial 4.0 Internacional ©iS

* This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. ©iS

to a more fundamental question about the existence of localizable particles.

The authors in (16) tackled the problem by demanding certain requirements from the states of a coordinate operator, which are used to localize elementary systems. Independently, among other operators the author in (17) gave as well a definition of an appropriate coordinate operator, which is equivalent to the one obtained by (16). Although the operator is unique for massive particles with arbitrary spin and for every massless system with spin 0, $1/2$, it has two major physical setbacks. First, eigenstates of the Newton-Wigner-Pryce (NWP) operator propagate superluminally and moreover the states are not covariant w.r.t. general Lorentz-transformations. Hence, there are two different opinions on this subject; either such localizable particles do not exist in a relativistic context or one can still make physical sense of the Newton-Wigner-Pryce operator and its respective states.

In this work, we take the second point of view and give, in our opinion, convincing arguments why the NWP states are of physical interest. In particular, the importance of these states relies on the fact that they can be used to calculate the probability amplitude of finding a particle at a certain spatial-position at time t . Moreover, the second quantization of this operator allows us to calculate the probability amplitude of finding k -particles at certain spatial-positions at time t .

In order to have a richer example of the NWP-space in a quantum field theoretical context (than the massive scalar field (Much)), we work with the free **massless** scalar field. In particular, we show that the localizability of particles, in the massless case, is not a lost cause. In general, the covariance group in the massless case is more than solely the Poincaré group, it is given by the conformal group. Hence, we investigate the covariant transformation property of NWP-states under conformal transformations. The importance therein has a physical reasoning. It is first of all done in order to prove that although the NWP-states are non-covariant w.r.t. to the whole conformal group there exist transformations of a subgroup of the conformal group that preserve their covariant character. Note that the dilatations, the spatial momentum and the pure rotations form indeed a subgroup of the conformal group that closes in the algebra. It is very often used in the context of quantum mechanics, see for example (2), (1) (and references therein). Furthermore, given two observers measuring, for a given covariant state, the probability of finding k -particles at k -spatial-positions. If these two observers are in two relatively to each other transformed frames, for example one is rotated, space-time translated or dilated to the other then they can agree on the observation of the probability of spatial positions, by using the covariant transformation rules.

Besides calculations of position-probability amplitudes of particles, the NWP-operator has an additional physical relevance. In this work we prove that by solely using the coordinate and relativistic momentum operator, one is able to define all generators of the conformal group. Hence, in some sense, the NWP-operator can be seen as essential for implementing relativistic symmetries.

Another motivation besides understanding localization to translate the conformal group into the coordinate space, is non-commutative (NC) geometry applied to quantum field theory (QFT). In particular the geometry that is quantized in NCQFT is the geometry of the NWP-space. Hence, in order to understand the outcome and the action of a deformation of QFT using, for example (see (14)) the conformal group, we need a better understanding of the commutative case.

The paper is organized as follows; Section two gives a quick introduction to the tools needed in this paper, where we define the Fock space of the free massless scalar field, the Newton-Wigner-Pryce

operator and the algebra of the Fourier-transformed ladder operators. In section four, the conformal algebra for a massless scalar field is transformed into the NWP-space. Section five investigates the covariance of the NWP-space under the Conformal group.

Conventions I.1. We use $d = n + 1$, for $n \in \mathbb{N}$ and the Greek letters are split into $\mu, \nu = 0, \dots, n$. Moreover, we use Latin letters for the spatial components which run from $1, \dots, n$ and we choose the following convention for the Minkowski scalar product of d -dimensional vectors, $a \cdot b = a_0 b^0 + a_k b^k = a_0 b^0 - \vec{a} \cdot \vec{b}$. Furthermore, we use the common symbol $\mathcal{S}(\mathbb{R}^n)$ for the Schwartz-space.

II | PRELIMINARIES

1 | Bosonic Fock-space and Fourier-transformation

We briefly define the Bosonic Fock space for a free scalar field ϕ with mass $m = 0$ on the $n + 1$ -dimensional Minkowski spacetime. In particular a particle with momentum $\mathbf{p} \in \mathbb{R}^n$ has the energy $\omega_{\mathbf{p}}$ given by $\omega_{\mathbf{p}} = |\mathbf{p}|$. Another quantity needed for the definition of the Fock-space is the Lorentz-invariant measure: $d^n \mu(\mathbf{p}) = d^n \mathbf{p} / (2\omega_{\mathbf{p}})$.

Definition The Bosonic Fock space \mathcal{H}^+ is defined as in (7; 21):

$$\mathcal{H}^+ = \bigoplus_{k=0}^{\infty} \mathcal{H}_k^+$$

where the k particle subspaces are given as

$$\begin{aligned} \mathcal{H}_k^+ &= \{\Psi_k : \partial V_+ \times \dots \times \partial V_+ \rightarrow \mathbb{C} \text{ symmetric} | \\ &\|\Psi_k\|^2 = \int d^n \mu(\mathbf{p}_1) \dots \int d^n \mu(\mathbf{p}_k) |\Psi_k(\mathbf{p}_1, \dots, \mathbf{p}_k)|^2 < \infty\}, \end{aligned}$$

with

$$\partial V_+ := \{p \in \mathbb{R}^{n+1} | p^2 = 0, p_0 > 0\}.$$

The particle annihilation and creation operators can be defined by their action on k -particle wave functions

$$\begin{aligned} (a_c(f)\Psi)_k(\mathbf{p}_1, \dots, \mathbf{p}_k) &= \sqrt{k+1} \int d^n \mu(\mathbf{p}) \overline{f(\mathbf{p})} \Psi_{k+1}(\mathbf{p}, \mathbf{p}_1, \dots, \mathbf{p}_k) \\ (a_c(f)^*\Psi)_k(\mathbf{p}, \mathbf{p}_1, \dots, \mathbf{p}_k) &= \begin{cases} 0, & k = 0 \\ \frac{1}{\sqrt{k}} \sum_{l=1}^k f(\mathbf{p}_l) \Psi_{k-1}(\mathbf{p}_1, \dots, \mathbf{p}_{l-1}, \mathbf{p}_{l+1}, \dots, \mathbf{p}_k), & k > 0 \end{cases} \end{aligned}$$

with $f \in \mathcal{H}_1$ and $\Psi_k \in \mathcal{H}_k^+$. The commutation relations of the smeared annihilation and creation operators $a_c(f), a_c(f)^*$ follow from their action on the functions Ψ and are given as follows

$$[a_c(f), a_c(g)^*] = \langle f, g \rangle = \int d^n \mu(\mathbf{p}) \overline{f(\mathbf{p})} g(\mathbf{p}), \quad [a_c(f), a_c(g)] = 0 = [a_c^*(f), a_c^*(g)].$$

The ladder operators with sharp momentum are introduced as operator valued distributions in the following

$$a_c(f) = \int d^n \mu(\mathbf{p}) \overline{f(\mathbf{p})} a_c(\mathbf{p}), \quad a_c(f)^* = \int d^n \mu(\mathbf{p}) f(\mathbf{p}) a_c^*(\mathbf{p}),$$

where the ladder operators with sharp momentum satisfy the well-known commutator relations

$$[a_c(\mathbf{p}), a_c(\mathbf{q})^*] = 2\omega_{\mathbf{p}}\delta^n(\mathbf{p} - \mathbf{q}), \quad [a_c(\mathbf{p}), a_c(\mathbf{q})] = 0 = [a_c^*(\mathbf{p}), a_c^*(\mathbf{q})].$$

In the following sections, we use the non-covariant normalization given by,

$$a(\mathbf{p}) = \frac{a_c(\mathbf{p})}{\sqrt{2\omega_{\mathbf{p}}}}, \quad a^*(\mathbf{p}) = \frac{a_c^*(\mathbf{p})}{\sqrt{2\omega_{\mathbf{p}}}}. \quad (1)$$

Next, we define the Fourier-transformation of the ladder operators that were given in the former expressions. These transformations are of physical importance in regards to the following sections.

Definition Fourier-transformation

In order to change from momentum space to the NWP-space we use explicit expressions for the Fourier-transformed creation and annihilation operators which are given by,

$$a(\mathbf{p}) = (2\pi)^{-n/2} \int d^n \mathbf{x} e^{i p_k x^k} \tilde{a}(\mathbf{x}), \quad a^*(\mathbf{p}) = (2\pi)^{-n/2} \int d^n \mathbf{x} e^{-i p_k x^k} \tilde{a}^*(\mathbf{x}).$$

The commutation relation between the ladder operators in momentum space gives us directly the relations for ladder operators of the NWP space,

$$\delta^n(\mathbf{p} - \mathbf{q}) = [a(\mathbf{p}), a^*(\mathbf{q})] = (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} e^{i p_k x^k} e^{-i q_k y^k} \underbrace{[\tilde{a}(\mathbf{x}), \tilde{a}^*(\mathbf{y})]}_{=\delta^n(\mathbf{x}-\mathbf{y})}.$$

Inverse Fourier-transformations of momentum space operators to the NWP-space ladder operators are given by,

$$\tilde{a}(\mathbf{x}) = (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-i p_k x^k} a(\mathbf{p}), \quad \tilde{a}^*(\mathbf{x}) = (2\pi)^{-n/2} \int d^n \mathbf{p} e^{i p_k x^k} a^*(\mathbf{p}). \quad (2)$$

2 | Massless NWP-operator

The (spatial)-QFT-position operator for a massive free scalar field is defined by the Newton-Wigner-Pryce operator, (17) and (16). On a one particle wave-function it acts as follows, (19, Chapter 3c, Equation 35)

$$(X_j \varphi)(\mathbf{p}) = -i \left(\frac{p_j}{2\omega_{m,\mathbf{p}}^2} + \frac{\partial}{\partial p^j} \right) \varphi(\mathbf{p}), \quad (3)$$

where $\omega_{m,\mathbf{p}}$ is the relativistic-energy for a particle with mass m . In order to prove that the NWP-operator is in the massless case equivalently represented as in the massive case, we can proceed in different ways. First, we can take Equation (3) and perform the massless limit. Since, the energy $\omega_{m,\mathbf{p}}$ is the only term that is affected by the limit, the massless NWP-operator is given by exchanging in Formula (3) the massive energy with $\omega_{\mathbf{p}}$. Therefore, the coordinate operator for a massless QFT has the same form as in the massive case. Another possible path to proceed is to define the coordinate operator as unitary equivalent to the second-quantized spatial-component of the relativistic momentum operator. In terms of the Fock space operators the relativistic momentum operator is given as,

$$P_\mu = \int d^n \mathbf{p} p_\mu a^*(\mathbf{p}) a(\mathbf{p}). \quad (4)$$

The unitary equivalence of the coordinate to the spatial momentum operator is given by the unitary map represented by the Fourier-transformation. In particular, it was proven that the NWP-operator can be represented as a self-adjoint operator on the domain, $\bigotimes_{i=1}^k \mathcal{S}(\mathbb{R}^n)$, with $\mathcal{S}(\mathbb{R}^n)$ denoting the Schwartz space, for details see (13) and (15). The explicit Fock-space representation of the spatial-coordinate operator is given by,

$$X_j = -i \int d^n \mathbf{p} a^*(\mathbf{p}) \frac{\partial}{\partial p^j} a(\mathbf{p}).$$

Since the spatial-momentum operator has the same form in the massless case as in the massive one, the NWP-operator, defined through the unitary equivalence, takes the same form as in the massive case. This stems from the fact that ladder operators do not explicitly depend on the mass. The commutation relations between the spatial-momentum operator and the NWP-operator are given by, see (23) or (15),

$$[X_j, P_k] = -i \eta_{jk} N, \quad (5)$$

where N is the particle-number operator represented in Fock-space as

$$N = \int d^n \mathbf{p} a^*(\mathbf{p}) a(\mathbf{p}). \quad (6)$$

Next, we want to give the physical interpretation and relevance of this operator. The eigenfunctions of the Newton-Wigner-Pryce operator, simultaneously representing the localized wave functions at time $x_0 = 0$, are given by, (19, Chapter 3, Equation 38)

$$\Psi_{\mathbf{x},0}(\mathbf{p}) = (2\pi)^{-n/2} e^{-i\mathbf{p}\cdot\mathbf{x}} (2\omega_p)^{1/2}.$$

Let a free massless scalar field be in a state $\Phi(\mathbf{p})$ at time $t = 0$. Then, the probability amplitude of finding a particle at the position \mathbf{x} is given by, (19, Chapter 3, Equation 44)

$$\langle \bar{\Psi}_{\mathbf{x}}, \Phi \rangle = \int d^n \mu(\mathbf{p}) \Psi_{\mathbf{x},0}(\mathbf{p}) \Phi(\mathbf{p}).$$

In order to extend this quantity to k -particles to calculate the probability amplitude for finding k -particle at positions $\mathbf{x}_1 \cdots \mathbf{x}_k$ at time $x_0 = x_{10} = \cdots = x_{k0}$, the following operator is introduced, (19, Chapter 7, Equation 99),

$$\phi_1(x) = \int d^n \mu(\mathbf{p}) \overline{\Psi_{\mathbf{x},x_0}(\mathbf{p})} a_c(\mathbf{p}) = (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-ip \cdot x} a(\mathbf{p}), \quad (7)$$

and we apply k -times ϕ_1 on $|\Phi^1\rangle$ and the vacuum as follows,

$$\Phi^1(x_1, \dots, x_k) = (k!)^{-1/2} \langle 0 | \phi_1(x_1) \cdots \phi_1(x_k) | \Phi^1 \rangle.$$

It is important to point out that the operator $\phi_1(x)$ at time $x^0 = 0$ is nothing else than the Fourier-transformed annihilation operator, see Equation (2). Moreover, the Newton-Wigner-Pryce operator has the following coordinate space representation (for proof see (Much, Lemma 3.1)),

$$X_j = \int d^n \mathbf{x} x_j \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}).$$

3| Constantly Used Integrals

To make this work self-contained, we give in this section a general formula for certain Fourier-transformed functions. In particular, these formulas can be found in (8, Chapter III, Section 2.6-2.8),

$$\begin{aligned}\tilde{P}^\lambda &= \int d^n \mathbf{p} |\tilde{p}|^{2\lambda} \exp\left(-i p_k z^k\right) \\ &= 2^{2\lambda+n} \pi^{\frac{1}{2}n} \frac{\Gamma(\lambda + \frac{1}{2}n)}{\Gamma(-\lambda)} (z_1^2 + \dots + z_n^2)^{-\lambda - \frac{1}{2}n},\end{aligned}\quad (8)$$

where Γ denotes the Gamma-function.

III | CONFORMAL GROUP IN THE NWP-SPACE

In this section we change the basis of the conformal group, which is given in terms of Fock-space operators in the momentum space, to the coordinate space. The base changes are mathematically well-defined on the domain $\bigotimes_{i=1}^k \mathcal{S}(\mathbb{R}^n)$. This conclusion follows from the fact that we use the Fourier-transformation for the base change and this transformation acts as a linear automorphism on the Schwartz-space. Hence, we conclude that the base change is well-defined.

However, an exception to the rule is given by the Lorentz generators generating boosts. They do not map the Schwartz-space to itself since the coefficient functions of the differential operators are not differentiable at $p = 0$. Hence, if one intends to calculate the adjoint action w.r.t. the Lorentz-boosts one has to choose the space of analytic functions on $S^2 \times \mathbb{R}$, more specifically of analytic sections, which is the light-cone without the tip, (see (4, Chapter 8.6)).¹

From a quantum field theoretical point of view, the conformal group is of great interest. In particular, the group is defined as the set of all conformal transformations. The definition of a conformal transformation is an invertible mapping $\mathbf{x}' \rightarrow \mathbf{x}$, leaving a d -dimensional metric g invariant. The invariance holds, modulo a scale factor, (3):

$$g'_{\mu\nu}(x') = F(x) g_{\mu\nu}(x). \quad (9)$$

Conformal mappings can be essentially summarized as Lorentz transformations, translations, dilations and special conformal transformations. The generators of these transformations are given by the operators $M_{\mu\nu}, P_\rho, D, K_\sigma$.

The conformal algebra is defined by the commutation relations of the generators and is given as follows:

$$[M_{\mu\nu}, M_{\rho\sigma}] = i(\eta_{\mu\sigma} M_{\nu\rho} + \eta_{\nu\rho} M_{\mu\sigma} - \eta_{\mu\rho} M_{\nu\sigma} - \eta_{\nu\sigma} M_{\mu\rho}), \quad (10)$$

$$[P_\rho, M_{\mu\nu}] = i(\eta_{\rho\mu} P_\nu - \eta_{\rho\nu} P_\mu), \quad [K_\rho, M_{\mu\nu}] = i(\eta_{\rho\mu} K_\nu - \eta_{\rho\nu} K_\mu), \quad (11)$$

$$[P_\rho, D] = iP_\rho, \quad [K_\rho, D] = -iK_\rho, \quad (12)$$

$$[P_\rho, K_\mu] = 2i(\eta_{\rho\mu} D - M_{\rho\mu}), \quad (13)$$

¹I am indebted to Prof. N. Dragon for this remark.

where all other commutators are equal to zero.

To transform the momentum and Lorentz operators in to the NWP-space we can take the massive expressions (see (Much)) and perform a massless limit, i.e. $m \rightarrow 0$. However, in order to make this work self-contained we calculate most expressions explicitly.

Lemma 1. *The zero component in the massless case is given by,*

$$P_0 = \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) (\tilde{\omega} * \tilde{a})(\mathbf{x}),$$

with function $\tilde{\omega}(x) := -\pi^{-\frac{n+1}{2}} \Gamma(\frac{n+1}{2}) |\mathbf{x}|^{-(n+1)}$ and $*$ denoting the convolution.

The massless spatial momentum operator takes the same form in coordinate space as in the massive case,

$$P_j = -i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) \frac{\partial}{\partial x^j} \tilde{a}(\mathbf{x}).$$

Demostración. For the proof we take the expression of the massless energy operator (see Equation (4)) and transform, by explicit Fourier-transformation, into the coordinate space,

$$\begin{aligned} P_0 &= \int d^n \mathbf{p} \omega_{\mathbf{p}} a^*(\mathbf{p}) a(\mathbf{p}) \\ &= (2\pi)^{-n} \int d^n \mathbf{p} |\mathbf{p}| \int d^n \mathbf{x} e^{i\mathbf{p} \cdot \mathbf{x}} \tilde{a}^*(\mathbf{x}) \int d^n \mathbf{y} e^{-i\mathbf{p} \cdot \mathbf{y}} \tilde{a}(\mathbf{y}) \\ &= (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} \left(\int d^n \mathbf{p} |\mathbf{p}| e^{i\mathbf{p} \cdot (\mathbf{x} - \mathbf{y})} \right) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\ &= -\pi^{-\frac{n+1}{2}} \Gamma(\frac{n+1}{2}) \iint d^n \mathbf{x} d^n \mathbf{y} \frac{1}{|\mathbf{x} - \mathbf{y}|^{n+1}} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}), \end{aligned}$$

where the Fourier-transformation can be found in (8, Chapter III, Section 2.6) or see Equation (8). The spatial momentum operator does not depend explicitly on the mass term, hence its representation in the massless is equivalent to the massive case, (see (Much)). \square

Other important expressions in the NWP-context are the velocity and the particle number operator. One way to calculate the velocity operator is by using the Heisenberg equation of motion,

$$[P_0, X_j] = -iV_j. \tag{14}$$

Since, we represented the relativistic energy and the NWP-operator in coordinate space, the calculation of the commutator gives us the velocity directly in coordinate space. In order to represent the particle number operator in the NWP-space we apply a Fourier-transformation on the expression in momentum space (see Equation (6)).

Lemma 2. *The velocity operator is given, in the massless case, as follows*

$$V_j = -i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) (\tilde{\omega}_j * \tilde{a})(\mathbf{x}), \tag{15}$$

with function $\tilde{\omega}_j(\mathbf{x}) := -\pi^{-\frac{n+1}{2}} \Gamma(\frac{n+1}{2}) |\mathbf{x}|^{-(n+1)} x_j$ and $*$ denoting the convolution.

Moreover, the particle number operator N is in the coordinate space given by,

$$N = \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}). \quad (16)$$

Demostración. We calculate the velocity by taking the commutator of the spatial coordinate space and the zero component of the momentum (see Heisenberg-Equation (14)), which is given in the former theorem, i.e.

$$\begin{aligned} [P_0, X_j] &= \iint d^n \mathbf{x} d^n \mathbf{z} z_j [\tilde{a}^*(\mathbf{x}) (\tilde{\omega} * \tilde{a})(\mathbf{x}), \tilde{a}^*(\mathbf{z}) \tilde{a}(\mathbf{z})] \\ &= \iiint d^n \mathbf{x} d^n \mathbf{y} d^n \mathbf{z} z_j \tilde{\omega}(\mathbf{x} - \mathbf{y}) \underbrace{[\tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}), \tilde{a}^*(\mathbf{z}) \tilde{a}(\mathbf{z})]}_{-\delta(\mathbf{x}-\mathbf{z}) \tilde{a}^*(\mathbf{z}) \tilde{a}(\mathbf{y}) + \delta(\mathbf{y}-\mathbf{z}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{z})} \\ &= - \iint d^n \mathbf{x} d^n \mathbf{y} (x - y)_j \tilde{\omega}(\mathbf{x} - \mathbf{y}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}). \end{aligned}$$

For the particle number operator we use the Fourier-transformation as in the proof of the former lemma,

$$\begin{aligned} N &= \int d^n \mathbf{p} \mathbf{p} a^*(\mathbf{p}) a(\mathbf{p}) \\ &= (2\pi)^{-n} \int d^n \mathbf{p} \int d^n \mathbf{x} e^{i p_k x^k} \tilde{a}^*(\mathbf{x}) \int d^n \mathbf{y} e^{-i p_l y^l} \tilde{a}(\mathbf{y}) \\ &= (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} \underbrace{\left(\int d^n \mathbf{p} e^{i p_k (x-y)^k} \right)}_{(2\pi)^n \delta(\mathbf{x}-\mathbf{y})} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}). \end{aligned}$$

□

Next, we turn our attention to the expressions of Lorentz generators. They are given as in the massive case by (see (11, Equation 3.54) and in this context see also (23, Appendix)),

$$M_{j0} = i \int d^n \mathbf{p} \mathbf{p} a^*(\mathbf{p}) \left(\frac{p_j}{2\omega_{\mathbf{p}}} - \omega_{\mathbf{p}} \frac{\partial}{\partial p^j} \right) a(\mathbf{p}), \quad (17)$$

$$M_{ik} = i \int d^n \mathbf{p} \mathbf{p} a^*(\mathbf{p}) \left(p_i \frac{\partial}{\partial p^k} - p_k \frac{\partial}{\partial p^i} \right) a(\mathbf{p}). \quad (18)$$

It is interesting and important to note that operators of boost and rotations can be given by the second quantization (denoted by $d\Gamma(\cdot)$ see (18, Chapter X.7)) of symmetric or skew-symmetric products of the momentum and coordinate operator.

Theorem 3. For the massless scalar field the generators of the proper orthochronous Lorentz-group \mathcal{L}_+^\uparrow can be written in terms of the second quantization of products of the NWP-operator with relativistic four-momentum as,

$$M_{0j} = \frac{1}{2} (d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)), \quad M_{ik} = d\Gamma(X_i P_k) - d\Gamma(X_k P_i). \quad (19)$$

Demostración. For the boosts we have,

$$\begin{aligned} M_{0j} &= \frac{1}{2} (d\Gamma(X_j P_0) + d\Gamma(P_0 X_j)) = \frac{1}{2} d\Gamma([X_j, P_0]) + d\Gamma(P_0 X_j) \\ &= \frac{i}{2} d\Gamma(V_j) + d\Gamma(P_0 X_j), \end{aligned}$$

where in the last lines we used the Heisenberg equation of motion. By comparing the last line with Formula (17) the proof follows. For rotations, the representation follows trivially. \square

The interesting fact about the former Theorem is that in principle one can define the generators of the Lorentz-group in a k -particle space by solely using the translation group and the NWP-operator. This resembles the proof of the Maxwell-equations by using the canonical commutation relations of Feynman (published by Dyson, (5)). In particular, the existence of the NWP-operator and the translation group are sufficient to build the Poincaré group, i.e. the group responsible for the implications of special relativity. Hence, the commutation relations with the addition of relativistic energy can be used to implement relativistic principles. This fact, is, in our opinion, an additional argument for the physical sense of the NWP-operator.

Next, we turn our attention to the explicit expressions of Lorentz generators in the coordinate space.

Lemma 4. *The boost generators of the Lorentz group expressed in the terms of ladder operators of the free massless scalar field are given in the coordinate space as*

$$M_{0j} = \frac{1}{2} \iint d^n \mathbf{x} d^n \mathbf{y} (x+y)_j \tilde{\omega}(\mathbf{x}-\mathbf{y}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}).$$

The operator of rotations takes the, from a quantum mechanical point of view, familiar and to the massive case equivalent form,

$$M_{ik} = i \int d^n \mathbf{x} \tilde{a}^*(\mathbf{x}) \left(x_i \frac{\partial}{\partial x^k} - x_k \frac{\partial}{\partial x^i} \right) \tilde{a}(\mathbf{x}).$$

Demostración. Since in the proof of (Much, Lemma 4.3) we did not explicitly encounter the mass when we calculated the representation of rotations in the coordinate space, we conclude that they have the same form. Next, we turn to the proof of the representation of Lorentz boosts in coordinate space. The first term is simply the massless velocity operator times $\frac{i}{2}$ (see Equation (19)). Hence, we focus here only on the second part,

$$\begin{aligned} & -i \int d^n \mathbf{p} a^*(\mathbf{p}) \omega_{\mathbf{p}} \frac{\partial}{\partial p^j} a(\mathbf{p}) \\ &= -i \int d^n \mathbf{p} \int d^n \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \omega_{\mathbf{p}} \frac{\partial}{\partial p^j} \int d^n \mathbf{y} e^{ip_s y^s} \tilde{a}(\mathbf{y}) \\ &= (2\pi)^{-n} \iint d^n \mathbf{x} d^n \mathbf{y} \left(\int d^n \mathbf{p} |\mathbf{p}| e^{-ip_k(x-y)^k} \right) y_j \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\ &= -\pi^{-\frac{n+1}{2}} \Gamma\left(\frac{n+1}{2}\right) \iint d^n \mathbf{x} d^n \mathbf{y} y_j \frac{1}{|\mathbf{x}-\mathbf{y}|^{n+1}} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \\ &= \iint d^n \mathbf{x} d^n \mathbf{y} y_j \tilde{\omega}(\mathbf{x}-\mathbf{y}) \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}), \end{aligned}$$

\square

In order to translate the special conformal and the dilatation operator into the coordinate space, we first write the momentum space representation, see (23). Moreover, we restrict this part to the physical space-time dimension four. The dilatation operator is given in momentum space as

$$D = -i \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(\frac{3}{2} + p^l \frac{\partial}{\partial p^l} \right) a(\mathbf{p}), \quad (20)$$

and for the special conformal operators we have,

$$K_0 = - \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(\frac{3}{4\omega_{\mathbf{p}}} + \frac{p^l}{\omega_{\mathbf{p}}} \frac{\partial}{\partial p^l} - \omega_{\mathbf{p}} \frac{\partial}{\partial p^l} \frac{\partial}{\partial p_l} \right) a(\mathbf{p}), \quad (21)$$

$$K_j = - \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(\frac{p_j}{4\omega_{\mathbf{p}}^2} + 3 \frac{\partial}{\partial p^j} + 2p^l \frac{\partial}{\partial p^l} \frac{\partial}{\partial p^j} - p_j \frac{\partial}{\partial p^l} \frac{\partial}{\partial p_l} \right) a(\mathbf{p}). \quad (22)$$

Lemma 5. *The four-dimensional dilatation operator is given in the coordinate representation as follows,*

$$D = i \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \left(\frac{3}{2} + x^l \frac{\partial}{\partial x^l} \right) \tilde{a}(\mathbf{x}). \quad (23)$$

Demostración. The Fourier-transformation is straight-forward in this case, e.g.

$$\begin{aligned} D &= -i \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(\frac{3}{2} + p^l \frac{\partial}{\partial p^l} \right) a(\mathbf{p}) \\ &= -i(2\pi)^{-3} \int d^3 \mathbf{p} \int d^3 \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \left(\frac{3}{2} + p^l \frac{\partial}{\partial p^l} \right) \int d^3 \mathbf{y} e^{ip_k y^k} \tilde{a}(\mathbf{y}) \\ &= -\frac{3i}{2} \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}) - i(2\pi)^{-3} \iint d^3 \mathbf{p} d^3 \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{y} \left(y^l \frac{\partial}{\partial y^l} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\ &= \frac{3i}{2} \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}) + i(2\pi)^{-3} \iint d^3 \mathbf{p} d^3 \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{y} e^{ip_k y^k} y^l \frac{\partial}{\partial y^l} \tilde{a}(\mathbf{y}) \\ &= i \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \left(\frac{3}{2} + x^l \frac{\partial}{\partial x^l} \right) \tilde{a}(\mathbf{x}). \end{aligned}$$

□

The outcome of the dilatation operator is interesting. In essence, it is equivalent to its momentum representation modulo a sign. This is what one might expect, since this operator performs dilatations in a complimentary manner, with regard to the momentum and coordinate space.

Next, we transform the special conformal operators into the coordinate space. From a calculational point of view these operators are the most difficult ones.

Lemma 6. *The zero component of the special conformal operators has the following form in coordinate space*

$$K_0 = \pi^{-2} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) |\mathbf{x} - \mathbf{y}|^{-2} \left(\frac{9}{8} - |\mathbf{x} - \mathbf{y}|^{-2} |\mathbf{y}|^2 + \frac{y^l}{2} \frac{\partial}{\partial y^l} \right) \tilde{a}(\mathbf{y}).$$

Demostración. In order to make this calculation more readable we separate the zero component into three parts as follows,

$$\begin{aligned} K_0 &= - \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(\frac{3}{4\omega_{\mathbf{p}}} + \frac{p^l}{\omega_{\mathbf{p}}} \frac{\partial}{\partial p^l} - \omega_{\mathbf{p}} \frac{\partial}{\partial p^l} \frac{\partial}{\partial p_l} \right) a(\mathbf{p}) \\ &=: K_0^1 + K_0^2 + K_0^3. \end{aligned}$$

Let us take a look at the first part,

$$\begin{aligned}
 K_0^1 &= -\frac{3}{4} \int d^3 \mathbf{p} a^*(\mathbf{p}) \omega_{\mathbf{p}}^{-1} a(\mathbf{p}) \\
 &= -\frac{3}{4} (2\pi)^{-3} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) \left(\int d^3 \mathbf{p} e^{-ip_k(x-y)^k} \omega_{\mathbf{p}}^{-1} \right) \tilde{a}(\mathbf{y}) \\
 &= -\frac{3}{8} \pi^{-2} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) |\mathbf{x} - \mathbf{y}|^{-2} \tilde{a}(\mathbf{y}).
 \end{aligned}$$

Next we turn our attention to the second term of the zero component of the special conformal operator,

$$\begin{aligned}
 K_0^2 &= - \int d^3 \mathbf{p} a^*(\mathbf{p}) \frac{p^l}{\omega_{\mathbf{p}}} \frac{\partial}{\partial p^l} a(\mathbf{p}) \\
 &= -(2\pi)^{-3} \int d^3 \mathbf{p} \int d^3 \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{y} \left(\frac{p^l}{\omega_{\mathbf{p}}} \frac{\partial}{\partial p^l} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\
 &= -(2\pi)^{-3} \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{p} \frac{1}{\omega_{\mathbf{p}}} e^{-ip_r x^r} \int d^3 \mathbf{y} \left(y^l \frac{\partial}{\partial y^l} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\
 &= (2\pi)^{-3} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) \left(\int d^3 \mathbf{p} \frac{1}{\omega_{\mathbf{p}}} e^{-ip_k(x-y)^k} \right) \left(3 + y^l \frac{\partial}{\partial y^l} \right) \tilde{a}(\mathbf{y}) \\
 &= (2)^{-1} \pi^{-2} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) (|\mathbf{x} - \mathbf{y}|^{-2}) \left(3 + y^l \frac{\partial}{\partial y^l} \right) \tilde{a}(\mathbf{y}),
 \end{aligned}$$

where in the last lines we performed a partial integration and used Formula (8). The last term of the zero component is given by

$$\begin{aligned}
 K_0^3 &= \int d^3 \mathbf{p} a^*(\mathbf{p}) \omega_{\mathbf{p}} \frac{\partial}{\partial p^l} \frac{\partial}{\partial p_l} a(\mathbf{p}) \\
 &= (2\pi)^{-3} \int d^3 \mathbf{p} \int d^3 \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{y} \left(\omega_{\mathbf{p}} \frac{\partial}{\partial p^l} \frac{\partial}{\partial p_l} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\
 &= (2\pi)^{-3} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) \left(\int d^3 \mathbf{p} \omega_{\mathbf{p}} e^{-ip_k(x-y)^k} \right) |\mathbf{y}|^2 \tilde{a}(\mathbf{y}) \\
 &= -(\pi)^{-2} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) (|\mathbf{x} - \mathbf{y}|^{-4}) |\mathbf{y}|^2 \tilde{a}(\mathbf{y}),
 \end{aligned}$$

where in the last lines we used the action of the differential operators w.r.t. \mathbf{p} on the exponential function and used as before Equation (8). □

Next, we give the base change of the spatial part of the special conformal operator.

Lemma 7. *The spatial components of the special conformal operators have the following form in coordinate space,*

$$\begin{aligned}
 K_j &= -\frac{i}{16} \pi^{-1} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) |\mathbf{x} - \mathbf{y}|^{-1} \frac{\partial}{\partial y^j} \tilde{a}(\mathbf{y}) \\
 &\quad + i \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \left(3x_j + 2x_j x^l \frac{\partial}{\partial x^l} - x_l x^l \frac{\partial}{\partial x^j} \right) \tilde{a}(\mathbf{x})
 \end{aligned} \tag{24}$$

Demostración. Let us first recall and define the expression of the special conformal operators in the

momentum space, i.e.

$$\begin{aligned} K_j &= - \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(\frac{p_j}{4\omega_{\mathbf{p}}^2} + 3 \frac{\partial}{\partial p^j} + 2p_l \frac{\partial}{\partial p_l} \frac{\partial}{\partial p^j} - p_j \frac{\partial}{\partial p^l} \frac{\partial}{\partial p_l} \right) a(\mathbf{p}) \\ &= K_j^1 - 3iX_j + K_j^2 + K_j^3. \end{aligned}$$

The second term in the spatial conformal operator is simply the coordinate operator times a constant. This fact was used in (22). Hence, the remaining terms of interest are K_j^1 , K_j^2 , K_j^3 . Let us start with first object,

$$\begin{aligned} K_j^1 &= -\frac{1}{4} \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(\frac{p_j}{\omega_{\mathbf{p}}^2} \right) a(\mathbf{p}) \\ &= -\frac{1}{4} (2\pi)^{-3} \int d^3 \mathbf{p} \int d^3 \mathbf{x} e^{-ip_r x^r} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{y} \left(\frac{p_j}{\omega_{\mathbf{p}}^2} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\ &= -\frac{i}{4} (2\pi)^{-3} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) \left(\int d^3 \mathbf{p} \frac{1}{\omega_{\mathbf{p}}^2} e^{-ip_k(x-y)^k} \right) \frac{\partial}{\partial y^j} \tilde{a}(\mathbf{y}) \\ &= -\frac{i}{16} \pi^{-1} \iint d^3 \mathbf{x} d^3 \mathbf{y} \tilde{a}^*(\mathbf{x}) |\mathbf{x} - \mathbf{y}|^{-1} \frac{\partial}{\partial y^j} \tilde{a}(\mathbf{y}), \end{aligned}$$

where in the last lines we expressed the momentum as a derivative, performed a partial integration and used the Fourier-transformation given in Formula (8) for $\lambda = -1$. A general term which includes, after taking the corresponding indices, the second and third operator of the spatial special conformal object is given by,

$$\begin{aligned} O_j &= \int d^3 \mathbf{p} a^*(\mathbf{p}) \left(p_r \frac{\partial}{\partial p_l} \frac{\partial}{\partial p^j} \right) a(\mathbf{p}) \\ &= (2\pi)^{-3} \int d^3 \mathbf{p} \int d^3 \mathbf{x} e^{-ip_s x^s} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{y} \left(p_r \frac{\partial}{\partial p_l} \frac{\partial}{\partial p^j} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\ &= i(2\pi)^{-3} \int d^3 \mathbf{p} \int d^3 \mathbf{x} e^{-ip_s x^s} \tilde{a}^*(\mathbf{x}) \int d^3 \mathbf{y} \left(y_j y^l \frac{\partial}{\partial y^r} e^{ip_k y^k} \right) \tilde{a}(\mathbf{y}) \\ &= -i \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \left(\eta_{rj} x^l + \eta_r^l x_j + x_j x^l \frac{\partial}{\partial x^r} \right) \tilde{a}(\mathbf{x}), \end{aligned}$$

where in the last lines we applied the derivatives to the exponential, expressed the coordinate y as a derivative and performed a partial integration. \square

Remark As for Lorentz generators, dilatation and special conformal operators can be written in terms of symmetric second-quantized products of the momentum and NWP-operator. For the dilatation operator it is straight forward,

$$D = \frac{1}{2} (d\Gamma(P_j X^j) + d\Gamma(X^j P_j)).$$

while the special-conformal operators are a bit more involved. For the zero component we have,

$$K_0 = -\frac{3}{4} d\Gamma(P_0^{-1}) - id\Gamma(V^l X_l) + d\Gamma(P_0 X_l X^l)$$

while the spatial components in terms of the NWP and the momentum operator read,

$$K_j = -\frac{1}{4}d\Gamma(P_0^{-1}V_j) - 3id\Gamma(X_j) + 2d\Gamma(P^lX_lX_j) - d\Gamma(P_jX_lX^l).$$

In this section we expressed all generators of the conformal group in terms of the ladder operators of the NWP space for $x_0 = 0$. However, the operator ϕ_1 (see Equation (7)) responsible for calculating the probability of finding k -particle at a certain position has an explicit time dependence. Hence, in order to connect our results with this time-dependency we give the following proposition.

Proposition 8. *The operator ϕ_1 that has the following form,*

$$\phi_1(x) = (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-ip \cdot x} a(\mathbf{p}),$$

can be obtained by performing a time-translation on the annihilation operator \tilde{a} of the NWP-space as follows

$$\phi_1(x^0, \mathbf{x}) = e^{ix^0 P_0} \tilde{a}(\mathbf{x}) e^{-ix^0 P_0}, \quad x^0 \in \mathbb{R}.$$

Demostración. By using the inverse Fourier-transformation of \tilde{a} and the transformation of the annihilation operator in momentum space under time-translation, the proof is completed, i.e.

$$\begin{aligned} e^{ix^0 P_0} \tilde{a}(\mathbf{x}) e^{-ix^0 P_0} &= (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-ip_k x^k} e^{ix^0 P_0} a(\mathbf{p}) e^{-iy^0 P_0} \\ &= (2\pi)^{-n/2} \int d^n \mathbf{p} e^{-ip_k x^k} e^{-ip_0 x^0} a(\mathbf{p}). \end{aligned}$$

□

Hence, we can simply time-translate all the operators of the conformal group in the NWP-space, by using the same time x^0 , in order to obtain the expressions for a general time and in terms of the operator ϕ_1 . Since the zero component of the momentum commutes with the spatial momentum and the rotations, we only have to calculate the time-translation for the remaining generators by using the algebra.

Lemma 9. *The dilatation transforms under time-translations as follows,*

$$e^{ix^0 P_0} D e^{-ix^0 P_0} = D - x^0 P_0,$$

where $x^0 \in \mathbb{R}$. Under time-translations the special conformal operator transforms as,

$$e^{ix^0 P_0} K_\mu e^{-ix^0 P_0} = K_\mu - 2x^0 (\eta_{0\mu} D - M_{0\mu}) - (x^0)^2 (P_\mu).$$

Demostración. The proof solely uses the conformal algebra (see Formulas (12) and (13)). For the dilatations we have,

$$e^{ix^0 P_0} D e^{-ix^0 P_0} = D - x^0 P_0 + \underbrace{\frac{i^2}{2!} (x^0)^2 [P_0, [P_0, D]]}_{=0} + \dots = D - x^0 P_0,$$

where in the last lines we used the Baker-Hausdorff formula. A more complex expression is the transformation of the special-conformal operator, i.e.

$$\begin{aligned} e^{ix^0 P_0} K_\mu e^{-ix^0 P_0} &= K_\mu + ix^0 [P_0, K_\mu] + \frac{i^2}{2!} (x^0)^2 [P_0, [P_0, K_\mu]] + \underbrace{\dots}_{=0} \\ &= K_\mu - 2x^0 (\eta_{0\mu} D - M_{0\mu}) - (x^0)^2 (\eta_{00} P_\mu), \end{aligned}$$

where we used the Baker-Campbell-Hausdorff formula and the specific commutator relations of the conformal algebra. □

Since, we have all expressions in terms of generators of the NWP-space we can equivalently give the conformal algebra expressed by operators $\phi_1^*(x_0, \mathbf{x})$ and $\phi_1(x_0, \mathbf{y})$. Note that neither the coordinate operator nor its respective eigenstates are covariant w.r.t. general Lorentz-transformations. Nevertheless, the generators of the conformal group obey the covariant transformation property independent of their representation. Hence, even though we use non-covariant eigenstates, operators of the conformal group represented in the NWP-space respect the Poincaré symmetry.

IV | CONFORMAL-TRANSFORMATIONS OF NWP-STATES

In this section we calculate the explicit adjoint action of the conformal group on the Fourier-transformed ladder operators. In some cases this is done by taking the massless limit of the massive theory.

The main physical motivation to calculate the adjoint action w.r.t. conformal transformations, in the coordinate space, is given by the fact that we use operator $\phi_1(x)$ to calculate the probability of localizing k -particles at k -spatial positions. Therefore, we intend to investigate under which subgroup of the conformal group the probability amplitudes of localization are covariant.

1 | Transformation under Translations

In order find the covariance group, we first define the unitary operator generating transformations of the orthochronous proper Poincaré group $\mathcal{P}_+^\uparrow = \mathcal{L}_+^\uparrow \ltimes \mathbb{R}^4$ denoted by $U(y, \Lambda)$. It transforms creation and annihilation operators in the following fashion, (19, Chapter 7),

$$U(y, \Lambda)a(\mathbf{p})U(y, \Lambda)^{-1} = \sqrt{\frac{\omega_{\Lambda\mathbf{p}}}{\omega_{\mathbf{p}}}} e^{-i(\Lambda p)_\mu y^\mu} a(\Lambda\mathbf{p}), \quad (y, \Lambda) \in \mathcal{P}_+^\uparrow, \quad (25)$$

$$U(y, \Lambda)a^*(\mathbf{p})U(y, \Lambda)^{-1} = \sqrt{\frac{\omega_{\Lambda\mathbf{p}}}{\omega_{\mathbf{p}}}} e^{i(\Lambda p)_\mu y^\mu} a^*(\Lambda\mathbf{p}), \quad (y, \Lambda) \in \mathcal{P}_+^\uparrow. \quad (26)$$

By using the former equations we first calculate the adjoint action of translations on the NWP-ladder operators.

Lemma 10. *The Fourier-transformed ladder operator \tilde{a} of the massless field transforms under translations as follows,*

$$U(y, \mathbb{I})\tilde{a}(\mathbf{x})U(y, \mathbb{I})^{-1} = c_y \left(\frac{1}{|\mathbf{x} + \mathbf{y}|^2 - (y^0)^2} \right)^2 * \tilde{a}(\mathbf{x} + \mathbf{y}), \quad (27)$$

where $c_y = \frac{iy^0}{\pi^2}$, $*$ denotes the convolution and $y \in \mathbb{R}^d$.

Demostración. The proof for spatial translations can be done as in the massive case (see (Much, Lemma 5.1)) since for this proof there is no explicit mass dependency. In particular they act accordingly

as spatial translations in coordinate space, i.e.

$$\begin{aligned} U(\mathbf{y}, \mathbb{I})\tilde{a}(\mathbf{x})U(\mathbf{y}, \mathbb{I})^{-1} &= (2\pi)^{3/2} \int d^3\mathbf{p} e^{-i\mathbf{p}_k x^k} U(\mathbf{y}, \mathbb{I})a(\mathbf{p})U(\mathbf{y}, \mathbb{I})^{-1} \\ &= (2\pi)^{-3/2} \int d^3\mathbf{p} e^{-i\mathbf{p}_k(x+y)^k} a(\mathbf{p}) = \tilde{a}(\mathbf{x} + \mathbf{y}). \end{aligned}$$

For time translations, we take the massless limit of the massive case (see (Much, Lemma 5)) where we had the integral,

$$(2\pi)^{-3} \int d^3\mathbf{p} e^{-i\mathbf{p}_k x^k} e^{-iy^0 \omega_{m,\mathbf{p}}} = \frac{iy^0}{2\pi^2} \frac{m^2 K_2(m\sqrt{|\mathbf{x}-\mathbf{z}|^2 - (y^0)^2})}{|\mathbf{x}-\mathbf{z}|^2 - (y^0)^2},$$

with $\omega_{m,\mathbf{p}}$ being the energy for a particle with mass m . By using the following limit,

$$\lim_{m \rightarrow 0} m^2 K_2(mf) = \frac{2}{f^2}, \quad (28)$$

the explicit form of the time translation can be given in the massless case as follows,

$$\begin{aligned} U(y^0, \mathbb{I})\tilde{a}(\mathbf{x})U(y^0, \mathbb{I})^{-1} &= e^{iy^0 P_0} \tilde{a}(\mathbf{x}) e^{-iy^0 P_0} \\ &= (2\pi)^{-3/2} \int d^3\mathbf{p} e^{-i\mathbf{p}_k x^k} e^{-iy^0 \omega_{\mathbf{p}}} a(\mathbf{p}) \\ &= (2\pi)^{-3} \int d^3\mathbf{z} \left(\int d^3\mathbf{p} e^{-i\mathbf{p}_k(x-z)^k} e^{-iy^0 \omega_{\mathbf{p}}} \right) \tilde{a}(\mathbf{z}) \\ &= \frac{iy^0}{\pi^2} \int d^3\mathbf{z} \left(\frac{1}{|\mathbf{x}-\mathbf{z}|^2 - (y^0)^2} \right)^2 \tilde{a}(\mathbf{z}). \end{aligned}$$

By taking the commutativity of the components of the momentum operator into account, we obtain the general transformation behavior under translations by using the former two results, i.e.

$$U(y, \mathbb{I})\tilde{a}(\mathbf{x})U(y, \mathbb{I})^{-1} = U(y^0, \mathbb{I})U(\mathbf{y}, \mathbb{I})\tilde{a}(\mathbf{x})U(\mathbf{y}, \mathbb{I})^{-1}U(y^0, \mathbb{I})^{-1}. \quad (29)$$

□

2 | Transformation under Pure Rotations

Before we turn to the Lorentz transformations in the massless case, we define the rotation group. In particular we denote matrices of the Lorentz group representing pure rotations as Λ_R . They are given by matrices of the following form,

$$\Lambda_R = \begin{pmatrix} 1 & 0 \\ 0 & R \end{pmatrix}, \quad R \in SO(3).$$

Lemma 11. *Under pure rotations the Fourier-transformed ladder operators \tilde{a} transforms in a covariant manner as follows,*

$$U(0, \Lambda_R)\tilde{a}(\mathbf{x})U(0, \Lambda_R)^{-1} = \tilde{a}(R\mathbf{x}).$$

Demostración. The proof is done analog to the proof of the former Lemma. However, we could take the results of (Much, Lemma 5.2) and perform the massless limit by using Equation (28). First, we

calculate the action of pure spatial rotations on the Fourier-transformed annihilation operator,

$$\begin{aligned}
 U(0, \Lambda_R) \tilde{a}(\mathbf{x}) U(0, \Lambda_R)^{-1} &= (2\pi)^{-3/2} \int d^3 \mathbf{p} e^{-i p_k x^k} U(0, \Lambda_R) a(\mathbf{p}) U(0, \Lambda_R)^{-1} \\
 &= (2\pi)^{-3/2} \int d^3 \mathbf{p} e^{-i p_k x^k} \sqrt{\frac{\Omega_{R\mathbf{p}}}{\Omega_{\mathbf{p}}}} a(R\mathbf{p}) \\
 &= (2\pi)^{-3/2} \int d^3 \mathbf{p} e^{-i p_k (R x)^k} a(\mathbf{p}),
 \end{aligned}$$

in the last lines we used the transformation behavior of the non-covariant momentum ladder operators (see (19, Chapter 7, Equation 57)) and the orthogonality of Λ_R for pure spatial rotations. \square

The NWP ladder operators transform non-covariantly w.r.t. the Lorentz-boosts. This is to be expected, since the NWP-operator is non-covariant w.r.t. the boosts. However, we see in the next theorem that the operator $\phi_1(x)$ is at least covariant w.r.t. space-time translations and pure rotations.

Theorem 12. *The NWP-ladder operators transform under space-time translations and pure rotations, i.e. with $(y, \Lambda_R) \in \mathcal{P}_+^\uparrow$, as follows,*

$$U(y, \Lambda_R) \tilde{a}(\mathbf{x}) U(y, \Lambda_R)^{-1} = \frac{iy^0}{\pi^2} \left(\frac{1}{|\mathbf{R}\mathbf{x} + \mathbf{y}|^2 - (y^0)^2} \right)^2 * \tilde{a}(\mathbf{R}\mathbf{x} + \mathbf{y}) \quad (30)$$

$$U(y, \Lambda_R) \tilde{a}^*(\mathbf{x}) U(y, \Lambda_R)^{-1} = -\frac{iy^0}{\pi^2} \left(\frac{1}{|\mathbf{R}\mathbf{x} + \mathbf{y}|^2 - (y^0)^2} \right)^2 * \tilde{a}^*(\mathbf{R}\mathbf{x} + \mathbf{y}). \quad (31)$$

Moreover, the operator used to calculate the probabilities, i.e. $\phi_1(x)$ transforms covariantly,

$$U(y, \Lambda_R) \phi_1(x) U(y, \Lambda_R)^{-1} = \phi_1(\Lambda_R x + y) \quad (y, \Lambda_R) \in \mathcal{P}_+^\uparrow. \quad (32)$$

Demostración. By using Lemmas 10 and 11 we deduce,

$$\begin{aligned}
 U(y, \Lambda_R) \tilde{a}(\mathbf{x}) U(y, \Lambda_R)^{-1} &= U(y, \mathbb{I}) U(0, \Lambda_R) \tilde{a}(\mathbf{x}) U(0, \Lambda_R)^{-1} U(y, \mathbb{I})^{-1} \\
 &= U(y, \mathbb{I}) \tilde{a}(\mathbf{R}\mathbf{x}) U(y, \mathbb{I})^{-1} \\
 &= \frac{iy^0}{\pi^2} \left(\frac{1}{|\mathbf{R}\mathbf{x} + \mathbf{y}|^2 - (y^0)^2} \right)^2 * \tilde{a}(\mathbf{R}\mathbf{x} + \mathbf{y}),
 \end{aligned}$$

where in the last lines we used the fact that a general Poincaré transformation can be split in first Lorentz-transformation and then space-time translation. An analog proof can be done for the creation operator \tilde{a}^* or we can use the fact that we have unitary operators generating the transformations and take the adjoint of the former equality.

In order to calculate the adjoint action of the operator $\phi_1(x)$, we use the fact that time translations commute with the unitary operator of the Poincaré group for pure rotations, i.e.

$$\begin{aligned}
 U(y, \Lambda_R) \phi_1(x) U(y, \Lambda_R)^{-1} &= U(y, \Lambda_R) U(x_0, \mathbb{I}) \tilde{a}(\mathbf{x}) U(x_0, \mathbb{I})^{-1} U(y, \Lambda_R)^{-1} \\
 &= U(x_0, \mathbb{I}) U(y, \Lambda_R) \tilde{a}(\mathbf{x}) U(y, \Lambda_R)^{-1} U(x_0, \mathbb{I})^{-1} \\
 &= U(x_0 + y_0, \mathbb{I}) U(\mathbf{y}, \mathbb{I}) \tilde{a}(\mathbf{R}\mathbf{x}) U(\mathbf{y}, \mathbb{I})^{-1} U(x_0 + y_0, \mathbb{I})^{-1} \\
 &= U(x_0 + y_0, \mathbb{I}) \tilde{a}(\mathbf{R}\mathbf{x} + \mathbf{y}) U(x_0 + y_0, \mathbb{I})^{-1} = \phi_1(\Lambda_R x + y).
 \end{aligned}$$

\square

3 | Transformation under Dilatations

Next, we investigate the adjoint action w.r.t. the dilatations. In order to calculate the adjoint action of dilatations on the NWP-space we first calculate the commutator of the dilatation operator with the coordinate operator. From that we can deduce the adjoint action of the dilatations acting on the coordinate ladder operators. This construction is analog to the one in (20, Chapter 9.4).

Lemma 13. *The adjoint action w.r.t. the dilatations acting on the NWP-operator is given as follows,*

$$e^{i\alpha D} X_j e^{-i\alpha D} = e^{-\alpha} X_j,$$

where in order to perform the calculations of the adjoint action the following commutator relation was used,

$$[D, X_j] = iX_j.$$

From the former equation it follows that $X_j e^{-i\alpha D} |\mathbf{x}\rangle$ is an eigenvector of X_j with eigenvalue $e^{-\alpha} x_j$. Therefore, one concludes that the transformation behavior of the operator $\tilde{a}(\mathbf{x})$ under dilatations is

$$e^{i\alpha D} \tilde{a}(\mathbf{x}) e^{-i\alpha D} = e^{\frac{3}{2}\alpha} \tilde{a}(e^\alpha \mathbf{x}).$$

Demostración. In order to proof the transformational behavior we first calculate the commutator of the dilatation operator with the NWP-operator,

$$\begin{aligned} [D, X_j] &= i \iint d^3 \mathbf{x} d^3 \mathbf{y} y_j [\tilde{a}^*(\mathbf{x}) \left(\frac{3}{2} + x^l \frac{\partial}{\partial x^l} \right) \tilde{a}(\mathbf{x}), \tilde{a}^*(\mathbf{y}) \tilde{a}(\mathbf{y})] \\ &= i \iint d^3 \mathbf{x} d^3 \mathbf{y} y_j x^l \underbrace{[\tilde{a}^*(\mathbf{x}) \frac{\partial}{\partial x^l} \tilde{a}(\mathbf{x}), \tilde{a}^*(\mathbf{y}) \tilde{a}(\mathbf{y})]}_{-\delta(\mathbf{x}-\mathbf{y}) \tilde{a}^*(\mathbf{y}) \frac{\partial}{\partial x^l} \tilde{a}(\mathbf{x}) + \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{y}) \frac{\partial}{\partial x^l} \delta(\mathbf{x}-\mathbf{y})} \\ &= -i \int d^3 x \left(x_j x^l \tilde{a}^*(\mathbf{x}) \frac{\partial}{\partial x^l} \tilde{a}(\mathbf{x}) + x_j \left(\frac{\partial}{\partial x^l} (x^l \tilde{a}^*(\mathbf{x})) \right) \tilde{a}(\mathbf{x}) \right) \\ &= -i \int d^3 x \left(x_j x^l \tilde{a}^*(\mathbf{x}) \frac{\partial}{\partial x^l} \tilde{a}(\mathbf{x}) - x^l \tilde{a}^*(\mathbf{x}) \left(\frac{\partial}{\partial x^l} (x_j \tilde{a}(\mathbf{x})) \right) \right) \\ &= i \int d^3 x x_j \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}), \end{aligned}$$

where in the last lines we used the vanishing commutator of the particle number operator with the coordinate operator. Moreover, we integrated over delta distributions and performed partial integrations.

From the former commutator relation we obtain for the adjoint action of the NWP-operator w.r.t. dilatations the following,

$$\begin{aligned} e^{i\alpha D} X_j e^{-i\alpha D} &= X_j + i\alpha [D, X_j] + \frac{i^2}{2!} \alpha^2 [D, [D, X_j]] + \dots \\ &= X_j + i^2 \alpha X_j + \frac{i^4}{2!} \alpha^2 X_j + \dots \\ &= \sum_{n=0}^{\infty} \frac{(i^2 \alpha)^n}{n!} X_j = e^{-\alpha} X_j, \end{aligned}$$

where in the last lines we used the commutator relation between the dilatation and the NWP-operator and

the Baker-Campbell-Hausdorff formula. As in (20, Chapter 9.4) we look at the following expression,

$$\begin{aligned} X_j e^{-i\alpha D} |\mathbf{x}\rangle &= e^{-i\alpha D} e^{i\alpha D} X_j e^{-i\alpha D} |\mathbf{x}\rangle \\ &= e^{-i\alpha D} e^{-\alpha} X_j |\mathbf{x}\rangle = e^{-\alpha} X_j e^{-i\alpha D} |\mathbf{x}\rangle. \end{aligned}$$

Therefore $X_j e^{-i\alpha D} |\mathbf{x}\rangle$ is an eigenvector of X_j for the eigenvalue $e^{-\alpha} X_j$, thus contained in the eigenspace $\mathcal{H}_{e^{-\alpha} X_j}$ of X_j . Hence, the transformation on the creation operator of the NWP-space has to have the following form,

$$e^{i\alpha D} \tilde{a}^*(\mathbf{x}) e^{-i\alpha D} = f(\alpha) \tilde{a}^*(e^{\alpha} \mathbf{x}), \quad (33)$$

where $f(\alpha)$ is a real-valued function of the transformation parameter α . To find the concrete form of the function f we calculate the adjoint action of the coordinate operator w.r.t. the dilatations but in the Fock representation of the NWP-space, i.e.

$$\begin{aligned} e^{i\alpha D} X_j e^{-i\alpha D} &= e^{i\alpha D} \int d^3 \mathbf{x} x_j \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}) e^{-i\alpha D} \\ &= \int d^3 \mathbf{x} x_j f(\alpha)^2 \tilde{a}^*(e^{\alpha} \mathbf{x}) \tilde{a}(e^{\alpha} \mathbf{x}) \\ &= e^{-\alpha} \int d^3 \mathbf{x} e^{-3\alpha} x_j f(\alpha)^2 \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}), \end{aligned}$$

where in order for the NWP-operator to transform correctly the function f has to be chosen as $f(\alpha) = e^{\frac{3}{2}\alpha}$. In order to solidify our approach we apply the adjoint action on the particle number operator in the coordinate space,

$$\begin{aligned} e^{i\alpha D} N e^{-i\alpha D} &= e^{i\alpha D} \int d^3 \mathbf{x} \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}) e^{-i\alpha D} \\ &= \int d^3 \mathbf{x} f(\alpha)^2 \tilde{a}^*(e^{\alpha} \mathbf{x}) \tilde{a}(e^{\alpha} \mathbf{x}) \\ &= \int d^3 \mathbf{x} e^{-3\alpha} f(\alpha)^2 \tilde{a}^*(\mathbf{x}) \tilde{a}(\mathbf{x}) = N, \end{aligned}$$

where in the last lines a variable substitution was performed and the commutator relation $[D, N] = 0$ is used to argue that the adjoint action leaves the particle number operator invariant and therefore the Ansatz (33) with explicit form of f gives us the right transformational behavior. \square

To calculate the adjoint action of dilatations on the ladder operators of the NWP-space we could have, as well, taken the well-known transformation property of the ladder operators in momentum space. After doing so we could have performed an inverse Fourier-transformation. However, one of the purposes of this work is to demonstrate to the reader that all calculations can as well be done in the NWP-space. In particular, the NWP-space is on an equal footing with the momentum space.

4 | Transformation under a Subgroup of the Conformal Group

Since we are interested in transformations of the time translated expression of the ladder operators we need to take the commutation relations between the momentum and dilatation operators into account.

Theorem 14. *The operator $\phi_1(x)$ that is responsible for calculating probabilities of finding k -particles in k -spatial positions transforms under dilatations as follows,*

$$e^{i\alpha D} \phi_1(x) e^{-i\alpha D} = e^{\frac{3}{2}\alpha} \phi_1(e^{\alpha} x).$$

Demostración. We can rewrite the operator $\phi_1(x)$ as

$$\phi_1(x) = U(x^0, \mathbb{I})\tilde{a}(\mathbf{x})U(-x^0, \mathbb{I}).$$

Since we know the adjoint action on the ladder operators in the NWP-space we can directly calculate the transformation on all expressions. First we calculate the adjoint action of the dilatations on the zero component of the momentum operator,

$$\begin{aligned} e^{i\alpha D} P_0 e^{-i\alpha D} &= -\pi^{-2}\Gamma(2) \iint d^3\mathbf{x}d^3\mathbf{y} \frac{1}{|\mathbf{x}-\mathbf{y}|^4} e^{i\alpha D} \tilde{a}^*(\mathbf{x}) e^{i\alpha D} e^{-i\alpha D} \tilde{a}(\mathbf{y}) e^{-i\alpha D} \\ &= -\pi^{-2}\Gamma(2) \iint d^3\mathbf{x}d^3\mathbf{y} \frac{1}{|\mathbf{x}-\mathbf{y}|^4} e^{3\alpha} \tilde{a}^*(e^\alpha\mathbf{x}) \tilde{a}(e^\alpha\mathbf{y}) \\ &= e^\alpha P_0, \end{aligned}$$

where in the last lines we used the explicit transformation given in Equation (33) and performed a variable substitution. Next, we apply the adjoint action of dilatations on this operator and use the fact that we are dealing with unitary transformations, i.e.

$$\begin{aligned} e^{i\alpha D} U(x^0, \mathbb{I})\tilde{a}(\mathbf{x})U(-x^0, \mathbb{I})e^{-i\alpha D} &= e^{\frac{3}{2}\alpha} U(e^\alpha x^0, \mathbb{I})\tilde{a}(e^\alpha\mathbf{x})U(-e^\alpha x^0, \mathbb{I}) \\ &= \frac{ie^{\frac{5}{2}\alpha} x^0}{2\pi^2} \int d^3\mathbf{z} \left(\frac{1}{|e^\alpha\mathbf{x}-\mathbf{z}|^2 - (e^\alpha x^0)^2} \right)^2 \tilde{a}(\mathbf{z}). \end{aligned}$$

□

By using the former lemmas we calculate the adjoint action of translations, rotations and dilatations on $\phi_1(x)$. This is in particular important since the operator $\phi_1(x)$ is used to calculate probability amplitudes of finding k -particles in k -spatial positions. Hence, by knowing the transformational behavior, two frames that are connected via translations, rotations and/or dilatations can calculate the same probability by taking the proper transformation into account.

Theorem 15. *The operator $\phi_1(x)$ transforms covariantly under space-time translations, pure rotations and dilatations, i.e.*

$$e^{i\alpha D} U(y, \Lambda_R)\phi_1(x)U(y, \Lambda_R)^{-1} e^{-i\alpha D} = e^{\frac{3}{2}\alpha} \phi_1(e^\alpha(\Lambda_R x + y)),$$

with $(y, \Lambda_R) \in \mathcal{P}_+^\uparrow$ and $\alpha \in \mathbb{R}$. Moreover, the explicit transformation of the Fourier-transformed operator \tilde{a} under the action of the subgroup $(y, \Lambda_R) \in \mathcal{P}_+^\uparrow$ is in coordinate space given as,

$$e^{i\alpha D} U(y, \Lambda_R)\tilde{a}(\mathbf{x})U(y, \Lambda_R)^{-1} e^{-i\alpha D} = c_{y'} \left(\frac{1}{|\mathbf{R}\mathbf{x} + \mathbf{y}|^2 - (y^0)^2} \right)^2 * \tilde{a}(e^\alpha(\mathbf{R}\mathbf{x} + \mathbf{y})),$$

where $c_{y'} := \frac{ie^{-\frac{3}{2}\alpha} y^0}{\pi^2}$, $*$ denotes the convolution and $y \in \mathbb{R}^4$.

Demostración. In order to prove the first part of the theorem we use the former Lemmas and the algebra of the Poincaré group. First, we examine how the operator $\phi_1(x)$ transforms under space-time

translations and pure rotations,

$$\begin{aligned}
 & e^{i\alpha D} U(y, \Lambda_R) \phi_1(x) U(y, \Lambda_R)^{-1} e^{-i\alpha D} \\
 &= e^{i\alpha D} \phi_1(\Lambda_R x + y) e^{-i\alpha D} \\
 &= e^{\frac{3}{2}\alpha} \phi_1(e^\alpha \Lambda_R x + e^\alpha y) \\
 &= e^{\frac{3}{2}\alpha} U(e^\alpha(x_0 + y_0), \mathbb{I}) \tilde{a}(e^\alpha(\mathbf{R}\mathbf{x} + \mathbf{y})) U(-e^\alpha(x_0 + y_0), \mathbb{I}) \\
 &= e^{\frac{3}{2}\alpha} c_{y''} \left(\frac{1}{|e^\alpha(\mathbf{R}\mathbf{x} + \mathbf{y})|^2 - (e^\alpha(x_0 + y_0))^2} \right)^2 * \tilde{a}(e^\alpha(\mathbf{R}\mathbf{x} + \mathbf{y})),
 \end{aligned}$$

where $c_{y''} := \frac{ie^\alpha(x_0 + y_0)}{\pi^2}$ and where for the calculation we used the transformation property given in Theorem 14, the dilatation action Equation (32) and the explicit time translation given in Equation (27). \square

V | CONCLUSION AND OUTLOOK

In this work we supported the point of view, that the NWP operator and its respective states have a physical meaning. In particular, we defined the second-quantized version of the NWP-operator for a massless scalar field and wrote the operator in its respective eigenbase. The meaning of the second-quantized operator is given by its respective eigenfunctions. They are used to calculate the probability amplitude of finding particles at certain spatial positions for a fixed time. Furthermore, by transforming the whole conformal group into the NWP-eigenbase we showed that this space can be treated and worked on at equal footing as the momentum space.

Although eigenstates of this operator are non-covariant w.r.t. the Poincaré group we showed indirectly that combinations of these states are covariant. This is proven by the fact that covariant operators remain covariant, independently of their respective representation. Hence, even though we work with non-covariant eigenstates, symmetries of the theory remain untouched. This is, in our opinion, another strong argument why the coordinate space, i.e. the NWP-space, is equally entitled from a physical point of view as the momentum space. However, the reason for the failure of Poincaré-covariance or covariance under the special conformal transformations is the fact that the states of the NW-operator are not covariant w.r.t. the before mentioned group. This as well implies the missing of a zero component of the NW-operator.

Moreover, we showed that the operator generating the NWP-space, i.e. $\phi_1(x)$ transforms in a covariant manner under the subgroup of space-time translations, rotations and dilatations of the conformal group, namely

$$e^{i\alpha D} U(y, \Lambda_R) \phi_1(x) U(y, \Lambda_R)^{-1} e^{-i\alpha D} = e^{\frac{3}{2}\alpha} \phi_1(e^\alpha(\Lambda_R x + y)).$$

Hence, given two observers that are related by space-time translations, rotations or/and dilatations, they both measure the same probability amplitude, by taking the respective covariant transformation into account.

Furthermore, we were able to supply an additional argument to point out the physical relevance of the NWP-operator. Namely, it is possible, even in a second-quantized context, to express the generators of the conformal group by using the NWP-operator and the relativistic momentum. In our opinion

this supplies a strong argument for the physical sense of the NWP-operator, since the conformal group can be written in terms of products of the coordinate and momentum operators.

Since we formulated the whole framework of a free massless scalar field, its respective transformations and symmetries into the NWP-space, the next step in progress is to understand localization from a non-commutative point of view. In particular, by imposing a space-space non-commutativity the subject of interest is to be able to calculate the probability of finding a particle at a certain mean value of a spatial position. In our opinion, the ground work for such investigations was presented in this paper.

I REFERENCIAS

- Chamon, C., Jackiw, R., Pi, S.-Y., and Santos, L. (2011). Conformal quantum mechanics as the CFT_1 dual to AdS_2 . *Phys. Lett.*, B701:503–507.
- de Alfaro, V., Fubini, S., and Furlan, G. (1976). Conformal invariance in quantum mechanics. *Il Nuovo Cimento A (1965-1970)*, 34(4):569–612.
- Di Francesco, P., Mathieu, P., and Sénéchal, D. (1997). *Conformal Field Theory*. Graduate Texts in Contemporary Physics. Springer.
- Dragon, N. (2012). *The Geometry of Special Relativity - A Concise Course*. SpringerBriefs in Physics. Springer Berlin Heidelberg.
- Dyson, F. J. (1990). Feynman's proof of the Maxwell equations. *Am.J.Phys.*, 58:209–211.
- Fleming, G. N. (1965a). Covariant Position Operators, Spin, and Locality. *Physical Review*, 137:B188–197.
- Fredenhagen, K. (2006). Quantum field theory, Lecture notes.
- Gel'fand, I. and Shilov, G. (1964). *Generalized functions*. Generalized Functions. Academic Press.
- Halvorson, H. (2000). Reeh-Schlieder Defeats Newton-Wigner: On alternative localization schemes in relativistic quantum field theory. *arXiv:quant-ph/0007060*.
- Herrmann, L. O. (2010). Localization in relativistic quantum theories.
- Itzykson, C. and Zuber, J. (2012). *Quantum Field Theory*. Dover Books on Physics. Dover Publications.
- Much, A. Remarks on QFT in the Coordinate Space. *arXiv:quant-ph/1609.01687*.
- Much, A. (2013). Quantum Spacetime from QM and QFT. *PhD thesis*.
- Much, A. (2014). Quantum Mechanical Effects from Deformation Theory. *Journal of Mathematical Physics*, 55(8):082303.
- Much, A. (2015). Relativistic Corrections to the Moyal-Weyl Spacetime. *Journal of Mathematical Physics*, 56(2):022301.
- Newton, T. D. and Wigner, E. P. (1949). Localized states for elementary systems. *Rev. Mod. Phys.*, 21:400–406.
- Pryce, M. H. L. (1948). The mass-centre in the restricted theory of relativity and its connexion with the quantum theory of elementary particles. *Proc. Royal Soc. London, Ser. A*, 195:62.
- Reed, M. and Simon, B. (1975). *Methods of Modern Mathematical Physics. 2. Fourier Analysis, Selfadjointness*. Gulf Professional Publishing.
- Schweber, S. (1961). *An Introduction to Relativistic Quantum Field Theory*. Harper & Row.
- Sextl, R. and Urbantke, H. (2012). *Relativity, Groups, Particles: Special Relativity and Relativistic Symmetry in Field and Particle Physics*. Springer Vienna.
- Sibold, K. (1993). Renormalization Theory, Lecture Notes.
- Sibold, K. and Burkhard, E. (2010). Conformal algebra on Fock space and conjugate pairs of operators. *Phys.Rev.*, D82:104012.
- Sibold, K. and Solard, G. (2009). Conjugate variables in quantum field theory: The basic case. *Phys.Rev.*, D80:124041.